

Projections in rotation algebras and theta functions

Florin P. Boca ^{*†‡}

e-mail fboca@stoilow.imar.ro, F.P.Boca@swansea.ac.uk

Abstract

For each $\alpha \in (0, 1)$, A_α denotes the universal C^* -algebra generated by two unitaries u and v , which fulfill the commutation relation $uv = e^{2\pi i \alpha} vu$. We consider the order four automorphism σ of A_α defined by $\sigma(u) = v$, $\sigma(v) = u^{-1}$ and describe a method for constructing projections in the fixed point algebra A_α^σ , using Rieffel's imprimitivity bimodules and Jacobi's theta functions. In the case $\alpha = q^{-1}$, $q \in \mathbf{Z}$, $q \geq 2$, we give explicit formulae for such projections and find some lower bounds for $\|u + u^* + v + v^*\|$.

The commutative algebra $C(\mathbf{T}^n)$ of continuous functions on the ordinary n -dimensional torus $\mathbf{T}^n = \{(z_1, \dots, z_n); |z_j| = 1\}$ is isomorphic to the universal C^* -algebra generated by n commuting unitary operators. A non-commutative n -torus A_α is the universal C^* -algebra generated by n unitaries $u_1, \dots, u_n$ subject to relations $u_j u_k = e^{2\pi i \alpha_{jk}} u_k u_j$, where $\alpha = (\alpha_{jk})_{1 \leq j, k \leq n}$ is a skew symmetric matrix with real entries. In some situations it is convenient to regard α as a real skew bilinear form on $\mathbf{Z}^n$ defined by $\alpha(e_j, e_k) = \alpha_{jk}$ and A_α as the twisted group C^* -algebra $C^*(\mathbf{Z}^n, \beta)$, where $\beta : \mathbf{Z}^n \times \mathbf{Z}^n \rightarrow \mathbf{T}$ is a 2-cocycle such that $\beta(x, y) \overline{\beta(y, x)} = e^{2\pi i \alpha(x, y)}$.

In this paper we only consider the case $n = 2$, when α is a real number and A_α is isomorphic to the crossed-product C^* -algebra $C(\mathbf{T}) \rtimes_\beta \mathbf{Z}$, where β is the automorphism of $C(\mathbf{T})$ defined by $\beta(\phi)(e^{2\pi i t}) = \phi(e^{2\pi i(t+\alpha)})$, $\phi \in C(\mathbf{T})$, $t \in \mathbf{R}$. The algebra A_α , called the rotation algebra by angle α , coincides with the universal C^* -algebra generated by two unitaries u and v which fulfill the commutation rule $uv = e^{2\pi i \alpha} vu$ and is endowed with the canonical faithful tracial state τ defined by $\tau\left(\sum_{m,n} a_{m,n} u^m v^n\right) = a_{0,0}$. The starting point in the study of rotation algebras is the existence of the Powers-Rieffel projections. They are projections $e_\alpha \in A_\alpha$ of trace α , which are of the form

$$e_\alpha = G(u)v + F(u) + \check{G}(u)v^{-1},$$

where $\check{G}(x) = G(-x)$, $x \in \mathbf{R}$ and F and G are some smooth functions on $\mathbf{R}$ (see [13]). The classical results of Pimsner and Voiculescu ([11],[12]) show that $\tau_*(K_0(A_\alpha)) = \mathbf{Z} + \mathbf{Z}\alpha = \mathbf{Z} + \mathbf{Z}\tau(e_\alpha)$; in particular, for any irrational numbers α_1 and α_2 , the rotation algebras A_{α_1} and A_{α_2} are isomorphic if and only if $\alpha_2 \pm \alpha_1 \in \mathbf{Z}$. The Powers-Rieffel projections play also a crucial rôle in the recent results ([6],[4]) on the structure of noncommutative tori.

The modular group $SL_2(\mathbf{Z})$ is a subgroup of the group $\text{Aut}(A_\alpha)$ of $*$ -automorphisms of A_α ; each matrix $X = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbf{Z})$ defines an automorphism σ_X of A_α , which acts on the canonical generators u and v of A_α as $\sigma_X(u) = u^a v^b$, $\sigma_X(v) = u^c v^d$. Throughout this paper, we will denote by $\sigma = \sigma\left(\begin{smallmatrix} 0 & 1 \\ -1 & 0 \end{smallmatrix}\right)$ the "Fourier transform" automorphism of A_α , acting on its generators by $\sigma(u) = v$ and $\sigma(v) = u^{-1}$. We also set

$$A_\alpha^\sigma = \{a \in A_\alpha; \sigma(a) = a\}$$

^{*}Research supported by an EPSRC Advanced Fellowship

[†]1991 *Mathematics Subject Classification*. 46L87, 14K25

[‡]The results of this paper were announced in the C^* -algebras meeting, Mathematisches Forschungsinstitut, Oberwolfach, February 1-7, 1998

and

$$H_\alpha^{(n)} = u^n + u^{-n} + v^n + v^{-n} \in A_\alpha^\sigma, \quad n \in \mathbf{Z}.$$

We notice that for any $\alpha \in \mathbf{R} \setminus \mathbf{Q}$, the C^* -algebra A_α^σ is generated only by the operators $H_\alpha = H_\alpha^{(1)}$ and $H_\alpha^{(2)}$ (see the appendix). The K-groups of A_α^σ were computed in the case $\alpha = \frac{p}{q} \in \mathbf{Q}$, $\gcd(p, q) = 1$ in [7], where it was shown that $K_1(A_\alpha^\sigma) = 0$ and $K_0(A_\alpha^\sigma) = \mathbf{Z}^9$ if $q \geq 5$.

The problem of characterizing the spectrum of the self-adjoint operator $H_\alpha = u + u^* + v + v^*$ (or more generally of $H_{\alpha, \lambda} = u + u^* + \frac{\lambda}{2}(v + v^*)$, $\lambda > 0$) in A_α is very important in the study of the quantum Hall effect ([2]). If $E_{\alpha, \lambda}$ denotes the spectral measure of $H_{\alpha, \lambda}$, then $\mu_{\alpha, \lambda} = \tau E_{\alpha, \lambda}$ is a measure with $\text{supp}(\mu_{\alpha, \lambda}) \subset [-2 - \lambda, 2 + \lambda]$ and since τ is faithful, its support coincides with the spectrum of $H_{\alpha, \lambda}$. Actually one can gather information on $\text{spec}(H_{\alpha, \lambda})$ from the K-theoretical properties of A_α . In this respect the results of Pimsner and Voiculescu ([12]) show that for any irrational α , any $\lambda \in \mathbf{R}$ and any t which belongs to a gap of $\text{spec}(H_{\alpha, \lambda})$, there exists an integer n such that

$$\mu_{\alpha, \lambda}(\text{spec}(H_{\alpha, \lambda}) \cap \chi_{(-\infty, t]}) = \{n\alpha\},$$

where $\{x\}$ denotes the fractional part of $x \in \mathbf{R}$. Therefore, knowing more about the projections of A_α^σ and about $\tau(\mathcal{P}(A_\alpha^\sigma))$, would presumably provide additional information on $\text{spec}(H_\alpha)$.

Another important feature of the automorphism σ is that it implements the Andre-Aubry duality. One easily checks that the set $\sigma_+(\alpha, \lambda) = \bigcup_{\theta} \sigma(\alpha, \lambda, \theta)$, as defined in [1], coincides with the spectrum of the operator $H_{\alpha, \lambda} \in A_\alpha$. Using the fact that σ is an automorphism of A_α and

$$\sigma(H_{\alpha, \lambda}) = \sigma\left(u + u^* + \frac{\lambda}{2}(v + v^*)\right) = v + v^* + \frac{\lambda}{2}(u + u^*) = \frac{\lambda}{2}\left(u + u^* + \frac{4}{\lambda} \cdot \frac{v + v^*}{2}\right) = \frac{\lambda}{2} H_{\alpha, \frac{4}{\lambda}},$$

it follows that the operators $H_{\alpha, \lambda}$ and $\frac{\lambda}{2} H_{\alpha, \frac{4}{\lambda}}$ have the same spectrum, hence we get a quick proof of the Andre-Aubry duality

$$\sigma_+(\alpha, \lambda) = \text{spec}(H_{\alpha, \lambda}) = \frac{\lambda}{2} \text{spec}(H_{\alpha, \frac{4}{\lambda}}) = \frac{\lambda}{2} \sigma_+(H_{\alpha, \frac{4}{\lambda}}).$$

The aim of this paper is to develop a method for constructing projections in the C^* -algebra A_α^σ . In the first section, we prove that for any $\alpha \in (0, 1)$, A_α^σ contains a projection e_α of trace α (the trace is the canonical trace on A_α). Although Rieffel's formalism ([14]) for constructing imprimitivity bimodules between twisted C^* -group algebras associated with lattices in abelian locally compact groups is quite useful, the nature of our projections is different from that of the Powers-Rieffel projections ([13]) used until now in the study of the rotation algebras. In fact, the projections we construct in A_α^σ are related to the classical Jacobi theta functions ([10], [9]). An important ingredient in the proof is the inequality

$$\vartheta(0, it)^2 < \vartheta(0, it) + \vartheta\left(\frac{1}{2}, it\right), \quad \forall t > \frac{1}{2},$$

which we prove in proposition 1.3 by making use of the infinite product factorization of theta functions; here we denote as usual

$$\vartheta(z, \tau) = \sum_m e^{\pi i m^2 \tau + 2\pi i m z}, \quad z \in \mathbf{C}, \tau \in \mathbf{H} = \{w \in \mathbf{C}; \text{Im}(w) > 0\}.$$

Another interesting fact related to this approach is that the classical transformation formula for $\vartheta(x, it)$ (see for example [10, p.33]) :

$$\vartheta\left(\frac{x}{it}, \frac{i}{t}\right) = \sqrt{t} e^{\frac{\pi x^2}{t}} \vartheta(x, it), \quad \forall t \in \mathbf{R}, t > 0,$$

arises as a mere consequence of Rieffel's trace formula ([14, thm.3.5]).

In section 2 we prove that if $p, q \in \mathbf{Z}$, $q \geq 2$, $0 < q\alpha - p < \frac{1}{2q}$ and p is a quadratic residue of q , then A_α^σ contains projections of trace $q\alpha - p$. If $0 < p - q\alpha < \frac{1}{2q}$ and $-p$ is a quadratic residue of q , then A_α^σ contains a projection of trace $p - q\alpha$.

The third section considers the case $\alpha = \frac{1}{q}$ $q \in \mathbf{Z}$, $q \geq 2$, in more detail. For example, we prove the following estimates for the norm of the operator H_α :

$$\begin{aligned} \|H_\alpha\| &\geq 4 e^{-\frac{\pi\alpha}{2}} \frac{\vartheta\left(\frac{1}{2}, \frac{i}{2\alpha}\right) \vartheta\left(\frac{i}{2}, \frac{i}{2\alpha}\right)}{\vartheta\left(0, \frac{i}{2\alpha}\right)^2} \quad \text{if } q \text{ is even} \\ \|H_\alpha\| &\geq 4 e^{-\frac{\pi\alpha}{2}} \frac{\vartheta\left(\frac{1}{2}, \frac{i}{2\alpha}\right) \vartheta\left(\frac{i}{2}, \frac{i}{2\alpha}\right) - 2 \vartheta^{odd}\left(\frac{1}{2}, \frac{i}{2\alpha}\right) \vartheta^{odd}\left(\frac{i}{2}, \frac{i}{2\alpha}\right)}{\vartheta\left(0, \frac{i}{2\alpha}\right)^2 - 2 \vartheta^{odd}\left(0, \frac{i}{2\alpha}\right)^2} \quad \text{if } q \text{ is odd,} \end{aligned}$$

where we set

$$\vartheta^{odd}(z, \tau) = \sum_{m \text{ odd}} e^{\pi i m^2 \tau + 2\pi i m z} = \vartheta(z, \tau) - \vartheta(2z, 4\tau), \quad z \in \mathbf{C}, \tau \in \mathbf{H}.$$

We also derive closed formulae for the projection e_α , expressing it as a sort of average of products of operator-valued $\vartheta_{a,b}$ functions (see proposition 3.1). Specializing for example to q even, it follows that the following identity holds for any integer $k \geq 1$ and $t_1, t_2 \in \mathbf{R}$:

$$\sum_{m,n=0}^{2k-1} e^{-\frac{2\pi i m n}{k}} \vartheta_{\frac{n}{2k}, \frac{m}{2}}(t_1, ik) \vartheta_{\frac{n}{2k}, \frac{m}{2}}(-t_1, ik) \vartheta_{\frac{m}{2k}, \frac{n}{2}}(t_2, ik) \vartheta_{\frac{m}{2k}, \frac{n}{2}}(-t_2, ik) = 2k \vartheta(t_1, ik)^2 \vartheta(t_2, ik)^2,$$

where we take as usual

$$\vartheta_{a,b}(z, \tau) = \sum_m e^{\pi i (m+a)^2 \tau + 2\pi i (m+a)(z+b)}, \quad z \in \mathbf{C}, \tau \in \mathbf{H}.$$

This should be compared with Riemann's identities for theta functions (see [10]).

1 Constructing projections of trace α in A_α^σ

We start by recalling the framework from [14]. Let M be an abelian locally compact group, let $\widehat{M}$ denote its topological dual and consider $G = M \times \widehat{M}$ equipped with the Haar-Plancherel measure. Let $\beta : G \times G \rightarrow \mathbf{T}$ be the Heisenberg bicharacter defined by

$$\beta((x_1, y_1), (x_2, y_2)) = \langle x_1, y_2 \rangle, \quad x_1, x_2 \in M, \quad y_1, y_2 \in \widehat{M},$$

where $\langle \cdot, \cdot \rangle : M \times \widehat{M} \rightarrow \mathbf{T}$ is the canonical pairing of M with $\widehat{M}$.

Set $\overline{\beta}(x, y) = \overline{\beta(x, y)}$ and $\beta^*(x, y) = \overline{\beta(y, x)} = \overline{\beta(y, x)}$, $x \in M$, $y \in \widehat{M}$. The formula

$$(\pi_{(x', x'')} f)(t) = \langle t, x'' \rangle f(t + x'), \quad t, x' \in M, \quad x'' \in \widehat{M}$$

defines a square-integrable projective unitary representation $\pi : G \rightarrow L^2(M)$ such that

$$\begin{aligned} \pi_x \pi_y &= \beta(x, y) \pi_{x+y}, \\ \pi_x \pi_y &= \beta \beta^*(x, y) \pi_y \pi_x, \\ (\pi_x)^* &= \beta(x, x) \pi_{-x}, \quad x, y \in G. \end{aligned}$$

If D is a lattice in G , we denote by $|G/D|$ its covolume and by $C^*(D, \beta)$ the C^* -algebra generated by π_w , $w \in D$. The subgroup

$$D^\perp = \{w \in G; \beta \beta^*(D, w) = 1\} = \{w \in G; \beta(x, w) = \beta(w, x), \forall x \in D\} \subset G$$

is a lattice in G . We equip D with the Haar measure which assigns mass $|G/D^\perp|^{-1} = |G/D|$ to each point and $D^\perp$ with the Haar measure assigning mass one to each point (see [14, p.278]). The twisted C^* -algebra $C^*(D, \beta)$ acts on the left on the space $\mathcal{S}(M)$ of Schwartz functions on M by

$$af = \int_D a(w) \pi_w f dw = |G/D| \sum_{w \in D} a(w) \pi_w f, \quad f \in \mathcal{S}(M), \quad a \in L^1(D, \beta). \quad (1.1)$$

Replacing as in [14, p.269] π_z by π_z^* , we regard $C^*(D^\perp, \beta)$ as being generated by π_z^* acting on the left on $\mathcal{S}(M)$. This action commutes with the left action of $C^*(D, \beta)$ for $\pi_w \pi_z = \pi_z \pi_w$, $w \in D$, $z \in D^\perp$. The opposite algebra of $C^*(D^\perp, \beta)$ is $C^*(D^\perp, \bar{\beta})$, which acts on the right on $\mathcal{S}(M)$ by

$$fb = \int_{D^\perp} b(z) (\pi_z^* f) dz = \sum_{z \in D^\perp} b(z) \pi_z^* f, \quad f \in \mathcal{S}(M), \quad b \in L^1(D^\perp, \bar{\beta}). \quad (1.2)$$

Moreover, $\mathcal{S}(M)$ becomes a $C^*(D, \beta) - C^*(D^\perp, \bar{\beta})$ equivalence bimodule with respect to the C^* -valued inner products $\langle \cdot, \cdot \rangle_D : \mathcal{S}(M) \times \mathcal{S}(M) \rightarrow C^*(D, \beta)$ and $\langle \cdot, \cdot \rangle_{D^\perp} : \mathcal{S}(M) \times \mathcal{S}(M) \rightarrow C^*(D^\perp, \bar{\beta})$ defined for any $f, g \in \mathcal{S}(M)$, $w = (w', w'') \in D$ and $z = (z', z'') \in D^\perp$ by

$$\langle f, g \rangle_D(w) = \langle f, \pi_w g \rangle_{L^2(M)} = \int_M f(s) \overline{g(s + w')} \overline{\langle s, w'' \rangle} ds, \quad (1.3)$$

$$\langle f, g \rangle_{D^\perp}(z) = \langle \pi_z g, f \rangle_{L^2(M)} = \int_M \overline{f(s)} g(s + z') \langle s, z'' \rangle ds. \quad (1.4)$$

If τ_D and $\tau_{D^\perp}$ are the canonical (normalized) traces on $C^*(D, \beta)$ and respectively on $C^*(D^\perp, \bar{\beta})$, then according to [14, thm.3.5] we have

$$\tau_D(\langle f, g \rangle_D) = |G/D| \tau_{D^\perp}(\langle g, f \rangle_{D^\perp}). \quad (1.5)$$

In this paper we are interested in the case when $M = \mathbf{R}^m \times F$, with F finite cyclic group. Since $\langle x, y \rangle = \mathbf{e}\left(\sum_{j=1}^m x_j y_j\right)$ if $x = (x_1, \dots, x_m)$, $y = (y_1, \dots, y_m) \in \mathbf{R}^m$ and $\langle [n]_q, [m]_q \rangle = \mathbf{e}\left(\frac{nm}{q}\right)$ if $[n]_q, [m]_q \in \mathbf{Z}_q$, where $\mathbf{e}(t) = \exp(2\pi i t)$ for all $t \in \mathbf{R}$, we shall identify M with $\widehat{M}$ such that $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in M = \widehat{M}$. Therefore $R(w_1, w_2) = (-w_2, w_1)$ defines a group automorphism of G . We notice that $R(D^\perp) = (RD)^\perp$.

The Fourier transform

$$(\mathcal{F}f)(s) = \int_M f(x) \overline{\langle x, s \rangle} dx, \quad f \in \mathcal{S}(M), \quad s \in M$$

extends to a unitary on $L^2(M)$ such that for all $g \in G$

$$\mathcal{F} \pi_g = \overline{\beta(g, g)} \pi_{Rg} \mathcal{F}. \quad (1.6)$$

Since $\beta(Rw, Rw) = \overline{\beta(w, w)}$, we see that for all $\xi_1, \xi_2 \in \mathcal{S}(M)$ and $w \in D$

$$\begin{aligned} \langle \mathcal{F} \xi_1, \mathcal{F} \xi_2 \rangle_D(Rw) &= \langle \mathcal{F} \xi_1, \pi_{Rw} \mathcal{F} \xi_2 \rangle_{L^2(M)} = \overline{\beta(w, w)} \langle \mathcal{F} \xi_1, \mathcal{F} \pi_w \xi_2 \rangle_{L^2(M)} \\ &= \overline{\beta(w, w)} \langle \xi_1, \pi_w \xi_2 \rangle_{L^2(M)} = \overline{\beta(w, w)} \langle \xi_1, \xi_2 \rangle_D(w), \end{aligned} \quad (1.7)$$

which yields for all $\xi_1, \xi_2, \xi_3 \in \mathcal{S}(M)$

$$\begin{aligned} \mathcal{F}(\langle \xi_1, \xi_2 \rangle_D \xi_3) &= \int_D \langle \xi_1, \xi_2 \rangle_D(w) \mathcal{F} \pi_w \xi_3 dw = \int_D \overline{\beta(w, w)} \langle \xi_1, \xi_2 \rangle_D(w) \pi_{Rw} \mathcal{F} \xi_3 dw \\ &= \int_D \langle \mathcal{F} \xi_1, \mathcal{F} \xi_2 \rangle_D(Rw) \pi_{Rw} \mathcal{F} \xi_3 dw = \langle \mathcal{F} \xi_1, \mathcal{F} \xi_2 \rangle_D(\mathcal{F} \xi_3). \end{aligned} \quad (1.8)$$

In a similar way, (1.4) and (1.6) yield for all $\xi_1, \xi_2 \in \mathcal{S}(M)$ and $z \in D^\perp$

$$\langle \mathcal{F}\xi_1, \mathcal{F}\xi_2 \rangle_{D^\perp}(Rz) = \langle \pi_{Rz} \mathcal{F}\xi_2, \mathcal{F}\xi_1 \rangle_{L^2(M)} = \beta(z, z) \langle \pi_z \xi_2, \xi_1 \rangle_{L^2(M)} = \beta(z, z) \langle \xi_1, \xi_2 \rangle_{D^\perp}(z),$$

and further on for all $\xi_1, \xi_2, \xi_3 \in \mathcal{S}(M)$

$$\begin{aligned} \mathcal{F}(\xi_1 \langle \xi_2, \xi_3 \rangle_{D^\perp}) &= \int_{D^\perp} \langle \xi_2, \xi_3 \rangle_{D^\perp}(z) \beta(z, z) \mathcal{F}\pi_{-z}\xi_1 dz = \int_{D^\perp} \langle \mathcal{F}\xi_2, \mathcal{F}\xi_3 \rangle_{D^\perp}(Rz) \pi_{Rz}^* \mathcal{F}\xi_1 dz \\ &= (\mathcal{F}\xi_1) \langle \mathcal{F}\xi_2, \mathcal{F}\xi_3 \rangle_{D^\perp}. \end{aligned} \quad (1.9)$$

According to [5], (1.8) and (1.9) show that $\mathbf{Z}_4 = \mathbf{Z}/4\mathbf{Z}$ acts on the imprimitivity bimodule $\mathcal{S}(M)$ by $u\xi = \mathcal{F}\xi$, inducing automorphisms $\sigma_D \in \text{Aut}(C^*(D, \beta))$ and $\sigma_{D^\perp} \in \text{Aut}(C^*(D^\perp, \beta))$ such that for all $\xi_1, \xi_2 \in \mathcal{S}(M)$, $a \in C^*(D, \beta)$, $b \in C^*(D^\perp, \beta)$

$$\sigma_D(\langle \xi_1, \xi_2 \rangle_D) = \langle \mathcal{F}\xi_1, \mathcal{F}\xi_2 \rangle_D, \quad (1.10)$$

$$\mathcal{F}(a\xi_1) = \sigma_D(a) (\mathcal{F}\xi_1), \quad (1.11)$$

$$\sigma_{D^\perp}(\langle \xi_1, \xi_2 \rangle_{D^\perp}) = \langle \mathcal{F}\xi_1, \mathcal{F}\xi_2 \rangle_{D^\perp}, \quad (1.12)$$

$$\mathcal{F}(\xi_1 b) = (\mathcal{F}\xi_1) \sigma_{D^\perp}(b). \quad (1.13)$$

To find σ_D , remark that (1.10) and (1.7) yield

$$\sigma_D(\langle \xi_1, \xi_2 \rangle_D) = \int_D \langle \mathcal{F}\xi_1, \mathcal{F}\xi_2 \rangle_D(Rw) \pi_{Rw} dw = \int_D \langle \xi_1, \xi_2 \rangle_D(w) \overline{\beta(w, w)} \pi_{Rw} dw.$$

On the other hand

$$\sigma_D(\langle \xi_1, \xi_2 \rangle_D) = \int_D \langle \xi_1, \xi_2 \rangle_D(w) \sigma_D(\pi_w) dw,$$

hence for all $w \in D$

$$\sigma_D(\pi_w) = \overline{\beta(w, w)} \pi_{Rw}. \quad (1.14)$$

A similar computation shows that for all $z \in D^\perp$

$$\sigma_{D^\perp}(\pi_z) = \overline{\beta(z, z)} \pi_{Rz}. \quad (1.15)$$

We notice that $\sigma_D^2(\pi_w) = \pi_{-w}$, $w \in D$, $\sigma_{D^\perp}(\pi_z) = \pi_{-z}$, $z \in D^\perp$, $\sigma_D^4 = \text{id}_{C^*(D, \beta)}$ and $\sigma_{D^\perp}^4 = \text{id}_{C^*(D^\perp, \beta)}$.

PROPOSITION 1.1 *Let $\alpha \in (0, 1)$ and $D = \mathbf{Z}\varepsilon_1 + \mathbf{Z}\varepsilon_2$ be a lattice in $G = \mathbf{R}^2$ such that $R\varepsilon_1 = \varepsilon_2$ (so $R\varepsilon_2 = -\varepsilon_1$) and $\beta(\varepsilon_j, \varepsilon_j) = 1$, $j = 1, 2$, $\beta\beta^*(\varepsilon_1, \varepsilon_2) = e^{2\pi i\alpha}$. Set $U_j = \pi_{\varepsilon_j}$, $j = 1, 2$ (hence $\sigma = \sigma_D$ is an automorphism of $A_\alpha = C^*(D, \beta)$ such that $\sigma(U_1) = U_2$, $\sigma(U_2) = U_1^{-1}$). Assume that there exists $\lambda \in \{\pm 1, \pm i\}$ and $f \in \mathcal{S}(\mathbf{R})$ such that $\mathcal{F}f = \lambda f$ and the element $a = \langle f, f \rangle_{D^\perp}$ is invertible in $C^*(D^\perp, \beta)$. Then $p = \langle fa^{-1/2}, fa^{-1/2} \rangle_D$ is a projection in A_α such that $\tau_D(p) = |G/D| = \alpha$ and $\sigma(p) = p$.*

Proof. The first part follows from [13], [14], so we only have to prove that $\sigma(p) = p$. Since $\mathcal{F}f = \lambda f$ and $R(D^\perp) = D^\perp$, (1.12) yields $\sigma_{D^\perp}(a) = a$, hence $\sigma_{D^\perp}(a^{-1/2}) = a^{-1/2}$. By (1.13) $\mathcal{F}(fa^{-1/2}) = \lambda fa^{-1/2}$ and applying (1.10) we get $\sigma_D(p) = p$. QED

Next, we choose $D = \mathbf{Z}\varepsilon_1 + \mathbf{Z}\varepsilon_2$, with $\varepsilon_1 = (\sqrt{\alpha}, 0)$, $\varepsilon_2 = (0, \sqrt{\alpha})$. The lattice D has covolume $|G/D| = \alpha$ in $G = \mathbf{R}^2$ and $RD = D$. Since $\beta\beta^*(\varepsilon_1, \varepsilon_2) = e^{2\pi i\alpha}$, the unitaries $U_j = \pi_{\varepsilon_j}$ fulfill $U_1 U_2 = e^{2\pi i\alpha} U_2 U_1$, hence $C^*(D, \beta)$ is canonically isomorphic to A_α . The automorphism $\sigma = \sigma_D$ acts on $A_\alpha = C^*(D, \beta)$ as $\sigma(\pi_{m_1\varepsilon_1 + m_2\varepsilon_2}) = e^{2\pi im_1 m_2 \alpha} \pi_{-m_2\varepsilon_1 + m_1\varepsilon_2}$, $m_1, m_2 \in \mathbf{Z}$, so $\sigma(U_1) = U_2$ and $\sigma(U_2) = U_1^{-1}$. Notice also that $\sigma^2(\pi_{m_1\varepsilon_1 + m_2\varepsilon_2}) = \pi_{-m_1\varepsilon_1 - m_2\varepsilon_2}$. The orthogonal lattice of D with respect to $\beta\beta^*$ is

$D^\perp = \mathbf{Z}\delta_1 + \mathbf{Z}\delta_2$, with $\delta_1 = (0, \frac{1}{\sqrt{\alpha}})$, $\delta_2 = (\frac{1}{\sqrt{\alpha}}, 0)$. It has covolume α^{-1} in $\mathbf{R}^2$ and $R(D^\perp) = D^\perp$. If we denote $V_1 = \pi_{\delta_1}^* = \pi_{-\delta_1}$ and $V_2 = \pi_{\delta_2}^* = \pi_{-\delta_2}$, then $\sigma_{D^\perp}(V_1) = V_2^{-1}$ and $\sigma_{D^\perp}(V_2) = V_1$.

Consider also the Schwartz function $f(s) = e^{-\pi s^2}$, $s \in \mathbf{R}$. Since $\mathcal{F}f = f$, proposition 1.1 shows that if $a = \langle f, f \rangle_{D^\perp}$ is invertible, then $\left\langle f \langle f, f \rangle_{D^\perp}^{-1/2}, f \langle f, f \rangle_{D^\perp}^{-1/2} \right\rangle_D$ is a projection of trace α in A_α^σ .

The following formula ([9, p.5]) will be used repeatedly throughout the paper

$$\int_{\mathbf{R}} e^{-2\pi s^2 + 2\pi a s} ds = \frac{1}{\sqrt{2}} e^{\frac{\pi a^2}{2}}, \quad \forall a \in \mathbf{C}. \quad (1.16)$$

Note also that for all $t > 0$

$$\int_{\mathbf{R}} e^{-\pi t s^2} ds = \frac{1}{\sqrt{t}}.$$

We make use of (1.4) and (1.16) to obtain

$$\begin{aligned} \langle f, f \rangle_{D^\perp}(m_1\delta_1 + m_2\delta_2) &= \langle f, f \rangle_{D^\perp} \left(\frac{m_2}{\sqrt{\alpha}}, \frac{m_1}{\sqrt{\alpha}} \right) = \int_{\mathbf{R}} \overline{f(s)} f \left(s + \frac{m_2}{\sqrt{\alpha}} \right) e^{\frac{2\pi i s m_1}{\sqrt{\alpha}}} ds \\ &= \int_{\mathbf{R}} e^{-2\pi s^2 - \frac{2\pi m_2 s}{\sqrt{\alpha}} + \frac{2\pi i m_1 s}{\sqrt{\alpha}} - \frac{\pi m_2^2}{\alpha}} ds = \frac{1}{\sqrt{2}} e^{-\frac{\pi(m_1^2 + m_2^2)}{2\alpha} - \frac{\pi i m_1 m_2}{\alpha}}, \end{aligned} \quad (1.17)$$

therefore

$$\begin{aligned} a = \langle f, f \rangle_{D^\perp} &= \sum_{z \in D^\perp} \langle f, f \rangle_{D^\perp}(z) \pi_z^* = \sum_{z \in D^\perp} \langle f, f \rangle_{D^\perp}(-z) \beta(z, z) \pi_z \\ &= \sum_{m_1, m_2} \langle f, f \rangle_{D^\perp}(-m_1\delta_1 - m_2\delta_2) e^{\frac{2\pi i m_1 m_2}{\alpha}} \pi_{m_1\delta_1 + m_2\delta_2} \\ &= \frac{1}{\sqrt{2}} \sum_{m_1, m_2} e^{-\frac{\pi(m_1^2 + m_2^2)}{2\alpha} + \frac{\pi i m_1 m_2}{\alpha}} V_1^{m_1} V_2^{m_2}. \end{aligned} \quad (1.18)$$

A direct computation based on (1.18) yields $2\tau_{D^\perp}(\langle f, f \rangle_{D^\perp}^2) = \vartheta(0, \frac{i}{\alpha})^2$. A computation similar to (1.17) yields $\sqrt{2} \langle f, f \rangle_D(m_1\varepsilon_1 + m_2\varepsilon_2) = e^{-\frac{\pi(m_1^2 + m_2^2)\alpha}{2} + \pi i m_1 m_2 \alpha}$, therefore

$$\sqrt{2} \langle f, f \rangle_D = |G/D| \sum_{w \in D} \langle f, f \rangle_D(w) \pi_w = \alpha \sum_{m_1, m_2} e^{-\frac{\pi(m_1^2 + m_2^2)\alpha}{2} + \pi i m_1 m_2 \alpha} U_2^{m_2} U_1^{m_1}, \quad (1.19)$$

which yields further $2\tau_D(\langle f, f \rangle_D^2) = \alpha^2 \vartheta(0, i\alpha)^2$. On the other hand, (1.5) yields for all $\phi, \psi \in \mathcal{S}(\mathbf{R})$

$$\tau_D(\langle \phi, \psi \rangle_D) = |G/D| \tau_{D^\perp}(\langle \psi, \phi \rangle_{D^\perp}) = \alpha \tau_{D^\perp}(\langle \psi, \phi \rangle_{D^\perp}), \quad (1.20)$$

therefore for all $f_1, f_2, f_3, f_4 \in \mathcal{S}(\mathbf{R})$

$$\begin{aligned} \tau_D(\langle f_1, f_2 \rangle_D \langle f_3, f_4 \rangle_D) &= \tau_D(\langle \langle f_1, f_2 \rangle_D f_3, f_4 \rangle_D) = \alpha \tau_{D^\perp}(\langle f_4, \langle f_1, f_2 \rangle_D f_3 \rangle_{D^\perp}) \\ &= \alpha \tau_{D^\perp}(\langle f_4, f_1 \rangle_{D^\perp} \langle f_2, f_3 \rangle_{D^\perp}). \end{aligned} \quad (1.21)$$

Taking $f_j = f$ in the previous equality we get

$$\vartheta(0, i\alpha) = \frac{1}{\sqrt{\alpha}} \vartheta\left(0, \frac{i}{\alpha}\right). \quad (1.22)$$

This is one of the modularity conditions satisfied by theta functions. Its appearance is not really surprising, for the Poisson summation formula plays an important rôle in the proof of (1.5) (and implicitly

of (1.20)). Actually we can do better by taking $f_a(s) = e^{-\pi(s+a)^2}$, $a \in \mathbf{R}$. A computation similar to the previous ones gives

$$\tau_D(2 \langle f_a, f \rangle_D \langle f, f_a \rangle_D) = \alpha^2 e^{-\pi a^2} \vartheta(-ia\sqrt{\alpha}, i\alpha) \vartheta(0, i\alpha),$$

$$\tau_{D^\perp}(2 \langle f_a, f_a \rangle_{D^\perp} \langle f, f \rangle_{D^\perp}) = \vartheta\left(\frac{a}{\sqrt{\alpha}}, \frac{i}{\alpha}\right) \vartheta\left(0, \frac{i}{\alpha}\right),$$

and using (1.21) and (1.22) :

$$\alpha e^{-\pi a^2} \vartheta(-ia\sqrt{\alpha}, i\alpha) \vartheta(0, i\alpha) = \vartheta\left(\frac{a}{\sqrt{\alpha}}, \frac{i}{\alpha}\right) \vartheta\left(0, \frac{i}{\alpha}\right) = \frac{1}{\sqrt{\alpha}} \vartheta\left(\frac{a}{\sqrt{\alpha}}, \frac{i}{\alpha}\right) \vartheta(0, i\alpha),$$

hence for all $a \in \mathbf{R}$ and $\alpha > 0$

$$\vartheta(-ia\sqrt{\alpha}, i\alpha) = \frac{1}{\sqrt{\alpha}} e^{\pi a^2} \vartheta\left(\frac{a}{\sqrt{\alpha}}, \frac{i}{\alpha}\right).$$

Taking $x = \frac{a}{\sqrt{\alpha}}$ and $t = \frac{1}{\alpha}$, we recover the following transformation formula for ϑ ([10, p.33])

$$\vartheta\left(\frac{x}{it}, \frac{i}{t}\right) = \sqrt{t} e^{\frac{\pi x^2}{t}} \vartheta(x, it), \quad \forall x \in \mathbf{R}, \quad \forall t > 0. \quad (1.23)$$

LEMMA 1.2 *The operator $X = \sum_m e^{-\pi m^2 \alpha_0} V_1^m$ is positive and invertible for all $\alpha_0 > 0$.*

Proof. As $X \in C^*(V_1) = C(\mathbf{T})$, we have $X(\lambda) = \sum_m e^{-\pi m^2 \alpha_0} \lambda^m$, $\lambda \in \mathbf{T}$, hence the spectrum of X coincides with $\vartheta([0, 1], i\alpha_0)$, where

$$\vartheta(z, \tau) = \sum_m e^{\pi i m^2 \tau + 2\pi i m z}, \quad z \in \mathbf{C}, \quad \tau \in \mathbf{H} = \{\zeta \in \mathbf{C}; \operatorname{Im} \zeta > 0\}$$

denotes the usual theta function ([10]). The operator X is self-adjoint for

$$X(e^{2\pi i t}) = 1 + 2 \sum_{m \geq 1} e^{-\pi m^2 \alpha_0} \cos(2\pi m t) \in \mathbf{R}, \quad \forall t \in \mathbf{R}.$$

On the other hand $X(1) = 1 + 2 \sum_{m \geq 1} e^{-\pi m^2 \alpha_0} > 0$, so we only have to show that $0 \notin \vartheta([0, 1], i\alpha_0)$.

This is true for $\frac{1+i\alpha_0}{2}$ is the only zero of the function $\theta_{\alpha_0}(z) = \vartheta(z, i\alpha_0)$ in the fundamental domain

$$\{z \in \mathbf{C}; 0 \leq \operatorname{Re} z \leq 1, 0 \leq \operatorname{Im} z \leq \alpha_0\}. \quad \text{QED}$$

Let $\alpha \in (0, \infty)$. We set

$$\alpha_m = e^{-\frac{\pi m^2}{2\alpha}},$$

$$\beta_m = \vartheta\left(-\frac{m}{2\alpha}, \frac{i}{2\alpha}\right), \quad m \in \mathbf{Z},$$

$$c(t) = \min_{x \in \mathbf{R}} \vartheta(x, it) > 0, \quad t > 0,$$

$$C(t) = \max_{x \in \mathbf{R}} \vartheta(x, it) = \vartheta(0, it) = \sum_n e^{-\pi n^2 t}, \quad t > 0,$$

$$\Phi_m(t) = \vartheta\left(-\frac{t}{\sqrt{\alpha}} + \frac{m}{2\alpha}, \frac{i}{2\alpha}\right), \quad m \in \mathbf{Z}.$$

We have $|\Phi_m(t)| \leq \sum_n e^{-\frac{\pi n^2}{2\alpha}} = \vartheta(0, \frac{i}{2\alpha}) = C(\frac{1}{2\alpha})$ for all $m \in \mathbf{Z}$ and $\alpha > 0$, hence the multiplication operator $D_m = M_{\Phi_m}$ is bounded on $L^2(\mathbf{R})$ and $\|D_m\| = \|\Phi_m\|_\infty \leq C(\frac{1}{2\alpha})$. On the other hand $\Phi_0(t) = \vartheta(-\frac{t}{\sqrt{\alpha}}, \frac{i}{2\alpha}) \in \mathbf{R}$ and $\Phi_0(t) \geq c(\frac{1}{2\alpha}) > 0$ for all $t \in \mathbf{R}$, hence D_0 is invertible and for all $m \in \mathbf{Z}$

$$\|D_0^{-1}D_m\| = \|M_{\frac{\Phi_m}{\Phi_0}}\| = \sup_{t \in \mathbf{R}} \left| \frac{\Phi_m(t)}{\Phi_0(t)} \right| \leq \frac{C(\frac{1}{2\alpha})}{c(\frac{1}{2\alpha})}. \quad (1.24)$$

We need more information on the behaviour of the function $\vartheta(\cdot, it)$ on $\mathbf{R}$ and prove the following

PROPOSITION 1.3 (i) For any $t > 0$

$$c(t) = \vartheta\left(\frac{1}{2}, it\right).$$

(ii) For any $t > 0.527$

$$\frac{C(t)(C(t) - 1)}{c(t)} < 1.$$

Proof. The proof relies on the infinite product expansion for theta functions (see for example [10, prop.14.1]), which says that for all $z \in \mathbf{C}$ and $\tau \in \mathbf{H}$ we have

$$\vartheta(z, \tau) = \prod_{m \geq 1} (1 - e^{2\pi i m \tau}) \prod_{m \geq 0} (1 + e^{(2m+1)\pi i \tau - 2\pi i z}) (1 + e^{(2m+1)\pi i \tau + 2\pi i z}). \quad (1.25)$$

Actually we will only use the "easy" fact that

$$\vartheta(z, \tau) = k_\tau \prod_{m \geq 0} (1 + e^{(2m+1)\pi i \tau - 2\pi i z}) (1 + e^{(2m+1)\pi i \tau + 2\pi i z}), \quad (1.26)$$

for some constant $k_\tau \neq 0$ which does not depend on z . To prove (i), remark that for any $r_m \geq 0$ and any $\rho \in \mathbf{T}$ we have

$$(1 + r_m \rho)(1 + r_m \bar{\rho}) \geq (1 - r_m)^2. \quad (1.27)$$

Taking $r = e^{-\pi t} \in (0, 1)$, $r_m = r^{2m+1}$ and $\rho = e^{2\pi i x}$, $x \in \mathbf{R}$, $t > 0$, we obtain from (1.26) and (1.27)

$$\frac{\vartheta(x, it)}{\vartheta(\frac{1}{2}, it)} = \prod_{m \geq 0} \frac{(1 + r_m \rho)(1 + r_m \bar{\rho})}{(1 - r_m)^2} \geq 1.$$

Since $\vartheta(x, it) > 0$ for all $x \in \mathbf{R}$, (i) follows.

To prove (ii), we fix $t > 0$ and denote $P = \sqrt{\frac{C(t)}{c(t)}} = \sqrt{\frac{\vartheta(0, it)}{\vartheta(\frac{1}{2}, it)}}$. Equality (1.26) yields :

$$\ln P = \sum_{m \geq 0} (\ln(1 + r^{2m+1}) - \ln(1 - r^{2m+1})).$$

The mean value theorem yields for any $\varepsilon \in (0, 1)$:

$$\frac{2\varepsilon}{1 + \varepsilon} < \ln(1 + \varepsilon) - \ln(1 - \varepsilon) < \frac{2\varepsilon}{1 - \varepsilon},$$

hence

$$\ln P < 2 \sum_{m \geq 0} \frac{r^{2m+1}}{1 - r^{2m+1}} < \frac{2r}{1 - r} \sum_{m \geq 0} r^{2m} = \frac{2r}{(1 - r)(1 - r^2)}$$

and therefore

$$\frac{C(t)}{c(t)} = P^2 < h(t) = \exp\left(\frac{4e^{-\pi t}}{(1 - e^{-\pi t})(1 - e^{-2\pi t})}\right) = \exp\left(\frac{4r}{(1 - r)(1 - r^2)}\right). \quad (1.28)$$

Using also

$$C(t) - 1 = \sum_{n \neq 0} e^{-\pi n^2 t} = 2 \sum_{n \geq 1} r^{n^2} < 2 \sum_{n \geq 0} r^{3n+1} = \frac{2r}{1-r^3},$$

we get

$$\frac{C(t)(C(t)-1)}{c(t)} < g(r) = \frac{2r}{1-r^3} \cdot \exp\left(\frac{4r}{(1-r)(1-r^2)}\right).$$

The derivative of

$$\phi(r) = \ln g(r) = \frac{4r}{(1-r)(1-r^3)} + \ln 2 + \ln\left(\frac{r}{1-r^3}\right), \quad r \in (0, 1),$$

is

$$\phi'(r) = \frac{4}{(1-r)(1-r^3)} + \frac{r(4r^2+r+1)}{(1-r)(1-r^3)^2} + \frac{1}{r} + \frac{3r^2}{1-r^3} > 0, \quad \forall r \in (0, 1),$$

therefore ϕ is monotonically increasing on $(0, 1)$. Moreover, the equation $\phi(r) = 0$ has a unique solution $r_0 \in (0, 1)$. If we set

$$\psi(x) = \phi\left(\frac{1}{x}\right) = \frac{4x^3}{(x-1)(x^3-1)} + \ln 2 + 2 \ln x - \ln(x^3-1), \quad x \in (1, \infty),$$

the equation $\psi(x) = 0$ has a unique solution $x_0 \approx 5.2254$ and ψ is monotonically decreasing on $(1, \infty)$. Therefore

$$C(t)(C(t)-1) < c(t), \quad \forall t > t_0 = \frac{1}{\pi} \log x_0 \approx 0.52633. \quad \text{QED}$$

PROPOSITION 1.4 *Let $\alpha \in (0, 0.948]$. Then $a_0 = \sum_m \alpha_m D_m V_2^m$ is a bounded invertible operator in $C^*(D^\perp, \beta) = A_{\frac{1}{\alpha}}$ and $a_0 = \sqrt{2} \langle f, f \rangle_{D^\perp}$.*

Proof. Since $\frac{1}{2\alpha} \geq 1 > 0.527$, (1.24) and the previous proposition yield

$$\sum_{m \neq 0} \alpha_m \|D_0^{-1} D_m V_2^m\| \leq \frac{C(\frac{1}{2\alpha})}{c(\frac{1}{2\alpha})} \sum_{m \neq 0} \alpha_m \leq \frac{C(\frac{1}{2\alpha}) (C(\frac{1}{2\alpha}) - 1)}{c(\frac{1}{2\alpha})} < 1.$$

This shows that $I + \sum_{m \neq 0} \alpha_m D_0^{-1} D_m V_2^m$ defines a bounded invertible operator, hence so is $a_0 = \sum_m \alpha_m D_m V_2^m$. The operators a_0 and $\sqrt{2} \langle f, f \rangle_{D^\perp} \in \mathcal{B}(L^2(\mathbf{R}))$ coincide. To see this, notice that for all $m_1, m_2 \in \mathbf{Z}$

$$(V_1^{m_1} V_2^{m_2} \phi)(s) = e^{-\frac{2\pi i m_1 s}{\sqrt{\alpha}}} \phi\left(s - \frac{m_2}{\sqrt{\alpha}}\right),$$

hence we may use (1.18) to obtain for all $\phi \in L^2(\mathbf{R})$ and $s \in \mathbf{R}$

$$\begin{aligned} \sqrt{2} (\langle f, f \rangle_{D^\perp}(\phi))(s) &= \sum_{m_1, m_2} e^{-\frac{\pi(m_1^2 + m_2^2)}{2\alpha} + \frac{\pi i m_1 m_2}{\alpha} - \frac{2\pi i m_1 s}{\sqrt{\alpha}}} \phi\left(s - \frac{m_2}{\sqrt{\alpha}}\right) \\ &= \left(\sum_{m_2} \alpha_{m_2} D_{m_2} V_2^{m_2} \phi \right)(s) = (a_0 \phi)(s). \end{aligned} \quad \text{QED}$$

COROLLARY 1.5 *For any $\alpha \in (0, 1)$, the rotation algebra A_α contains a projection $e = e_\alpha$ of trace α such that $\sigma(e) = e$.*

REMARKS 1.6 (i) For all $m \in \mathbf{Z}$ we have $D_m V_2 = V_2 D_{m+2}$.

(ii) If α is irrational, the rotation algebra $A_{\frac{1}{\alpha}} = C^*(D^\perp, \beta)$ is simple, thus is isomorphic to the C^* -algebra $C^*(W_1, W_2) \subset \mathcal{B}(\ell^2(\mathbf{Z}))$ generated by the unitaries $W_1 \xi_k = \xi_{k+1}$, $W_2 \xi_k = e^{\frac{2\pi i k}{\alpha}} \xi_k$, where ξ_k is an orthonormal basis of $\ell^2(\mathbf{Z})$. If $\rho = e^{\frac{2\pi i}{\alpha}}$, then the matrix coefficients of $a = \langle f, f \rangle_{D^\perp}$ in this representation of $A_{\frac{1}{\alpha}}$ are

$$\begin{aligned} \langle a \xi_k, \xi_l \rangle &= \frac{1}{\sqrt{2}} \sum_{m_1, m_2} \rho^{\frac{m_1 m_2}{2}} \alpha_{m_1} \alpha_{m_2} \langle W_1^{m_1} W_2^{m_2} \xi_k, \xi_l \rangle = \frac{1}{\sqrt{2}} \sum_{m_1, m_2} \rho^{\frac{m_1 m_2}{2} + m_2 k} \alpha_{m_1} \alpha_{m_2} \langle \xi_{k+m_1}, \xi_l \rangle \\ &= \frac{1}{\sqrt{2}} \alpha_{l-k} \sum_{m_2} \rho^{\frac{(k+l)m_2}{2}} \alpha_{m_2} = \frac{1}{\sqrt{2}} \alpha_{l-k} \vartheta \left(\frac{k+l}{2\alpha}, \frac{i}{2\alpha} \right) = \frac{1}{\sqrt{2}} \alpha_{l-k} \beta_{l+k}, \quad k, l \in \mathbf{Z}, \end{aligned}$$

therefore $a\sqrt{2} = \sum_m \alpha_m D_m U^m$, where $U \xi_k = \xi_{k+1}$, $D_m \xi_k = \beta_{2k-m} \xi_k$, $m, k \in \mathbf{Z}$. The diagonal operators are bounded and invertible because $0 < c \leq \beta_n \leq C$ for all $n \in \mathbf{Z}$.

2 Existence of projections of trace $q\alpha - p$ and $p - q\alpha$

Let $q \in \mathbf{N}^*$, $q \geq 2$ and $p \in \mathbf{Z}$ such that $0 < \gamma = \alpha - \frac{p}{q} \leq \frac{1}{2}$ and there exists $p_0 \in \mathbf{Z}$ such that $p = p_0^2 \bmod q$.

We choose $M = \mathbf{R} \times \mathbf{Z}_q$ and $D = \mathbf{Z}\varepsilon_1 + \mathbf{Z}\varepsilon_2 \subset G = M \times \widehat{M}$, with

$$\varepsilon_1 = (\sqrt{\gamma}, [p_0]_q, 0, [0]_q) \quad \text{and} \quad \varepsilon_2 = (0, [0]_q, \sqrt{\gamma}, [p_0]_q).$$

Then D is a lattice in G and $[0, \sqrt{\gamma}) \times \mathbf{Z}_q \times [0, \sqrt{\gamma}) \times \mathbf{Z}_q$ a fundamental domain for G/D . Since $\mathbf{Z}_q$ is endowed with the Haar-Plancherel measure (which assigns to each point mass $q^{-1/2}$), we get $|G/D| = q\gamma = q\alpha - p$. An easy computation gives $D^\perp = \mathbf{Z}\delta_1 + \mathbf{Z}\delta_2$ with

$$\delta_1 = \left(0, [0]_q, \frac{1}{q\sqrt{\gamma}}, [\bar{p}]_q \right) \quad \text{and} \quad \delta_2 = \left(\frac{1}{q\sqrt{\gamma}}, [\bar{p}]_q, 0, [0]_q \right),$$

where $\bar{p} \in \mathbf{Z}$ is such that $p_0 \bar{p} = -1 \bmod q$. Set $V_j = \pi_{\delta_j}^* = \pi_{-\delta_j}$, $j = 1, 2$. For any $\phi \in \mathcal{S}(\mathbf{R})$, we consider $\phi_1, \phi_2 \in \mathcal{S}(M)$ defined by

$$\phi_1(s, [n]_q) = \phi(s)$$

$$\phi_2(s, [n]_q) = \sqrt{q} \delta_{[0]_q, [n]_q} \phi(s), \quad s \in \mathbf{R}, \quad [n]_q \in \mathbf{Z}_q,$$

where $\delta_{a,b}$, $a, b \in \mathbf{Z}_q$ denotes Kronecker's symbol. Denote also $\delta_a(b) = \delta_{a,b}$, $a, b \in \mathbf{Z}_q$. Notice that if $\mathcal{F}\phi = \phi$ on $\mathbf{R}$, then $\mathcal{F}(\phi_1 + \phi_2) = \phi_1 + \phi_2$ on M because $\mathcal{F}(1 + \sqrt{q} \delta_{[0]_q}) = 1 + \sqrt{q} \delta_{[0]_q}$ (again, it is essential that the measure on $\mathbf{Z}_q$ is the Haar-Plancherel one). If we set

$$b_{m_1, m_2} = \int_{\mathbf{R}} \overline{\phi(s)} \phi\left(s + \frac{m_2}{q\sqrt{\gamma}}\right) e^{\frac{2\pi i s m_1}{q\sqrt{\gamma}}} ds, \quad m_1, m_2 \in \mathbf{Z},$$

then

$$\begin{aligned}
\langle \phi_1, \phi_1 \rangle_{D^\perp} (m_1 \delta_1 + m_2 \delta_2) &= \langle \phi_1, \phi_1 \rangle_{D^\perp} \left(\frac{m_2}{q\sqrt{\gamma}}, [m_2 \bar{p}]_q, \frac{m_1}{q\sqrt{\gamma}}, [m_1 \bar{p}]_q \right) \\
&= \int_{\mathbf{R} \times \mathbf{Z}_q} \overline{\phi_1(s, [n]_q)} \phi_1 \left(s + \frac{m_2}{q\sqrt{\gamma}}, [n + m_2 \bar{p}]_q \right) e^{\frac{2\pi i s m_1}{q\sqrt{\gamma}} + \frac{2\pi i n m_1 \bar{p}}{q}} ds d[n]_q \\
&= \frac{1}{\sqrt{q}} \sum_{n \in \mathbf{Z}_q} e^{\frac{2\pi i n m_1 \bar{p}}{q}} b_{m_1, m_2} = \begin{cases} 0, & \text{if } q \nmid m_1 \\ \sqrt{q} b_{m_1, m_2}, & \text{if } q \mid m_1 \end{cases}, \\
\langle \phi_1, \phi_2 \rangle_{D^\perp} (m_1 \delta_1 + m_2 \delta_2) &= \int_{\mathbf{R} \times \mathbf{Z}_q} \overline{\phi_1(s, [n]_q)} \phi_2 \left(s + \frac{m_2}{q\sqrt{\gamma}}, [n + m_2 \bar{p}]_q \right) \\
&\quad \cdot e^{\frac{2\pi i s m_1}{q\sqrt{\gamma}} + \frac{2\pi i n m_1 \bar{p}}{q}} ds d[n]_q = e^{-\frac{2\pi i m_1 m_2 \bar{p}^2}{q}} b_{m_1, m_2}, \\
\langle \phi_2, \phi_1 \rangle_{D^\perp} (m_1 \delta_1 + m_2 \delta_2) &= b_{m_1, m_2}, \\
\langle \phi_2, \phi_2 \rangle_{D^\perp} (m_1 \delta_1 + m_2 \delta_2) &= \begin{cases} 0, & \text{if } q \nmid m_2 \\ \sqrt{q} b_{m_1, m_2}, & \text{if } q \mid m_2 \end{cases}
\end{aligned} \tag{2.1}$$

and consequently for all $m_1, m_2 \in \mathbf{Z}$

$$\langle \phi_1 + \phi_2, \phi_1 + \phi_2 \rangle_{D^\perp} (m_1 \delta_1 + m_2 \delta_2) = b_{m_1, m_2} c_{m_1, m_2},$$

where

$$c_{m_1, m_2} = \begin{cases} 1 + e^{-\frac{2\pi i m_1 m_2 \bar{p}^2}{q}}, & \text{if } q \nmid m_1 \text{ and } q \nmid m_2 \\ 2 + \sqrt{q}, & \text{if } q \nmid m_1, q \mid m_2 \text{ or } q \mid m_1, q \nmid m_2 \\ 2 + 2\sqrt{q}, & \text{if } q \mid m_1 \text{ and } q \mid m_2 \end{cases}.$$

If $\phi(s) = e^{-\pi s^2}$, set $\tilde{\phi} = \phi_1 + \phi_2$. By the computations from section 1 (with $q\sqrt{\gamma}$ instead of $\sqrt{\alpha}$) we get for all $m_1, m_2 \in \mathbf{Z}$

$$b_{m_1, m_2} = \frac{1}{\sqrt{2}} e^{-\frac{\pi(m_1^2 + m_2^2)}{2q^2\gamma} - \frac{\pi i m_1 m_2}{q^2\gamma}},$$

and

$$\begin{aligned}
a\sqrt{2} &= \sqrt{2} \langle \tilde{\phi}, \tilde{\phi} \rangle_{D^\perp} = \sum_{m_1, m_2} b_{m_1, m_2} c_{m_1, m_2} \pi_{m_1 \delta_1 + m_2 \delta_2}^* \\
&= \sqrt{2} \sum_{m_1, m_2} b_{-m_1, -m_2} c_{-m_1, -m_2} \beta(-m_2 \delta_2, -m_1 \delta_1) \pi_{m_1 \delta_1 + m_2 \delta_2} \\
&= \sqrt{2} \sum_{m_1, m_2} b_{-m_1, -m_2} c_{-m_1, -m_2} V_2^{m_2} V_1^{m_1} = \sqrt{2} \sum_{m_1, m_2} b_{m_1, m_2} c_{m_1, m_2} V_2^{m_2} V_1^{m_1} \\
&= \sum_{m_1, m_2} c_{m_1, m_2} e^{-\frac{\pi(m_1^2 + m_2^2)}{2q^2\gamma} - \frac{\pi i m_1 m_2}{q^2\gamma}} V_2^{m_2} V_1^{m_1} \\
&= \sum_{m_2} e^{-\frac{\pi m_2^2}{2q^2\gamma}} \left(\sum_{m_1} c_{m_1, m_2} e^{\frac{2\pi i m_1 m_2 \bar{p}^2}{q} + \frac{\pi i m_1 m_2}{q^2\gamma} - \frac{\pi m_1^2}{2q^2\gamma}} V_1^{m_1} \right) V_2^{m_2}.
\end{aligned}$$

We set $\alpha_m = e^{-\frac{\pi m^2}{2q^2\gamma}}$, $m \in \mathbf{Z}$ and

$$D_n = \sum_m c_{m, n} \alpha_m e^{\frac{2\pi i m n}{q} \left(\frac{1}{2q\gamma} + \bar{p}^2 \right)} V_1^m, \quad n \in \mathbf{Z}.$$

For all $f \in L^2(\mathbf{R} \times \mathbf{Z}_q)$

$$(V_1^m f)(x, [k]_q) = e^{-\frac{2\pi i m x}{q\sqrt{\gamma}} - \frac{2\pi i m k \bar{p}}{q}} f(x, [k]_q),$$

hence D_n is the multiplication operator M_{Φ_n} on $L^2(\mathbf{R} \times \mathbf{Z}_q)$, with $\Phi_n \in L^\infty(\mathbf{R} \times \mathbf{Z}_q)$ given by

$$\Phi_n(x, [k]_q) = \sum_m c_{m,n} \alpha_m e^{\frac{2\pi i m n}{q} \left(\frac{1}{2q\gamma} + \bar{p}^2 \right) - \frac{2\pi i m x}{q\sqrt{\gamma}} - \frac{2\pi i m k \bar{p}}{q}}$$

and $a\sqrt{2} = \sum_n \alpha_n D_n V_2^n$. We have

$$\begin{aligned} \Phi_0(x, [k]_q) &= \sum_m c_{m,0} \alpha_m e^{-\frac{2\pi i m x}{q\sqrt{\gamma}} - \frac{2\pi i m k \bar{p}}{q}} \\ &= (2 + 2\sqrt{q}) \sum_l \alpha_{ql} e^{-\frac{2\pi i l x}{\sqrt{\gamma}}} + (2 + \sqrt{q}) \sum_{q|l} \alpha_m e^{-\frac{2\pi i m x}{q\sqrt{\gamma}} - \frac{2\pi i m k \bar{p}}{q}} \\ &= \sqrt{q} \vartheta\left(-\frac{x}{\sqrt{\gamma}}, \frac{i}{2\gamma}\right) + (2 + \sqrt{q}) \vartheta\left(-\frac{x}{q\sqrt{\gamma}} - \frac{k\bar{p}}{q}, \frac{i}{2q^2\gamma}\right) \\ &\geq (2 + \sqrt{q}) c\left(\frac{1}{2q^2\gamma}\right) + \sqrt{q} c\left(\frac{1}{2\gamma}\right) > 0, \end{aligned} \tag{2.2}$$

where $c(t) = \min_{x \in \mathbf{R}} \vartheta(x, it) = \vartheta(\frac{1}{2}, it)$, $t > 0$ and

$$\begin{aligned} |\Phi_n(x, [n]_q)| &\leq \sum_m |c_{m,n}| \alpha_m = \sum_l c_{ql,n} \alpha_{ql} + \sum_{q|l} |c_{m,n}| \alpha_m \\ &\leq (2 + 2\sqrt{q}) \vartheta\left(0, \frac{i}{2\gamma}\right) + (2 + \sqrt{q}) \left(\vartheta\left(0, \frac{i}{2q^2\gamma}\right) - \vartheta\left(0, \frac{i}{2\gamma}\right) \right) \\ &= (2 + \sqrt{q}) C\left(\frac{1}{2q^2\gamma}\right) + \sqrt{q} C\left(\frac{1}{2\gamma}\right), \end{aligned} \tag{2.3}$$

where $C(t) = \max_{x \in \mathbf{R}} \vartheta(x, it) = \vartheta(0, it)$, $t > 0$. We combine (2.2) and (2.3) to obtain for all $n \in \mathbf{Z}$

$$\left\| \frac{\Phi_n}{\Phi_0} \right\|_\infty = \sup \left\{ \left| \frac{\Phi_n(x, [k]_q)}{\Phi_0(x, [k]_q)} \right|; (x, [k]_q) \in \mathbf{R} \times \mathbf{Z}_q \right\} \leq \frac{(2 + \sqrt{q}) C\left(\frac{1}{2q^2\gamma}\right) + \sqrt{q} C\left(\frac{1}{2\gamma}\right)}{(2 + \sqrt{q}) c\left(\frac{1}{2q^2\gamma}\right) + \sqrt{q} c\left(\frac{1}{2\gamma}\right)}. \tag{2.4}$$

The function h from (1.28) is monotonically decreasing on $(0, \infty)$ and $C(t) < h(t) c(t)$ for all $t > 0$, therefore (2.4) yields for all $n \in \mathbf{Z}$

$$\left\| \frac{\Phi_n}{\Phi_0} \right\|_\infty \leq \max \left\{ h\left(\frac{1}{2q^2\gamma}\right), h\left(\frac{1}{2\gamma}\right) \right\} = h\left(\frac{1}{2q^2\gamma}\right),$$

and

$$S = \sum_{n \neq 0} \alpha_n \left\| \frac{\Phi_n}{\Phi_0} \right\|_\infty \leq 2 h\left(\frac{1}{2q^2\gamma}\right) \sum_{n \geq 1} \alpha_n.$$

But

$$\sum_{n \geq 1} \alpha_n = \sum_{n \geq 1} e^{-\frac{\pi n^2}{2q^2\gamma}} < \sum_{n \geq 0} e^{-\frac{\pi(3n+1)}{2q^2\gamma}} = \frac{e^{-\frac{\pi}{2q^2\gamma}}}{1 - e^{-\frac{3\pi}{2q^2\gamma}}},$$

therefore

$$S \leq 2h\left(\frac{1}{2q^2\gamma}\right) \cdot \frac{e^{-\frac{\pi}{2q^2\gamma}}}{1 - e^{-\frac{3\pi}{2q^2\gamma}}} = g(r),$$

where $r = e^{-\frac{\pi}{2q^2\gamma}}$ and g is as in the proof of proposition 1.7. We have already shown thereby that $g(r) < 1$ if $\frac{1}{2q^2\gamma} > 0.527$, hence in this case $I + \sum_{n \neq 0} \alpha_n D_0^{-1} D_n V_2^n$ is bounded and invertible on $L^2(\mathbf{R} \times \mathbf{Z}_q)$, hence $\sqrt{2} \langle \tilde{\phi}, \tilde{\phi} \rangle_{D^\perp} = \sum_n \alpha_n D_n V_2^n$ is invertible and we have proved the following

PROPOSITION 2.1 *Let $\alpha \in (0, 1)$. If $q \geq 2$ and p are integers such that $q \geq 2$, $0 < \alpha - \frac{p}{q} < \frac{0.948}{q^2}$ and there exists $p_0 \in \mathbf{Z}$ such that $p = p_0^2 \bmod q$, then a is invertible and A_α^σ contains a projection of trace $q\alpha - p$.*

One proves in a similar way that if $0 < \gamma' = \frac{p}{q} - \alpha < \frac{0.948}{q^2}$ and there exists $p_1 \in \mathbf{Z}$ such that $p = -p_1^2 \bmod q$, then a is invertible, hence A_α^σ contains a projection of trace $p - q\alpha$. The computations are as above, starting with $\varepsilon_1 = (0, [0]_q, \sqrt{\gamma'}, [p_1]_q)$ and $\varepsilon_2 = (\sqrt{\gamma'}, [p_1]_q, 0, [0]_q)$.

3 More on the case $\alpha = q^{-1}$, $q \in \mathbf{Z}$, $q \geq 2$

In this section we will focus mainly on the case $\alpha = \frac{1}{q}$, $q \in \mathbf{Z}$, $q \geq 2$, obtaining lower bounds for the norm of $H_\alpha = \pi_{\varepsilon_1} + \pi_{\varepsilon_1}^* + \pi_{\varepsilon_2} + \pi_{\varepsilon_2}^*$ and closed formulae for the projection constructed in corollary 1.5 (notation is as in section 1).

In the beginning we will assume that $\alpha \in (0, 1)$ is such that the operator $\langle f, f \rangle_{D^\perp}$ from (1.18) is invertible (for example $0 < \alpha < 0.948$), hence $e = \left\langle f \langle f, f \rangle_{D^\perp}^{-\frac{1}{2}}, f \langle f, f \rangle_{D^\perp}^{-\frac{1}{2}} \right\rangle_D$ is a projection of trace α in A_α^σ . Then, according to [13], $\Theta : eA_\alpha e \simeq eC^*(D, \beta)e \rightarrow C^*(D^\perp, \bar{\beta}) \simeq A_{\frac{1}{\alpha}}^\perp$ defined by

$$\Theta(exe) = \langle f, f \rangle_{D^\perp}^{-\frac{1}{2}} \langle f, xf \rangle_{D^\perp} \langle f, f \rangle_{D^\perp}^{-\frac{1}{2}}, \quad x \in C^*(D, \beta)$$

is a *-isomorphism. Direct computations based on (1.4) and (1.16) yield

$$\begin{aligned} \langle f, \pi_{\pm \varepsilon_1} f \rangle_{D^\perp} (m_1 \delta_1 + m_2 \delta_2) &= \langle f, \pi_{\pm \varepsilon_1} f \rangle_{D^\perp} \left(\frac{m_2}{\sqrt{\alpha}}, \frac{m_1}{\sqrt{\alpha}} \right) = \int_{\mathbf{R}} \overline{f(s)} f \left(s + \frac{m_2}{\sqrt{\alpha}} \pm \sqrt{\alpha} \right) e^{\frac{2\pi i s m_1}{\sqrt{\alpha}}} ds \\ &= \int_{\mathbf{R}} e^{-\pi s^2 - \pi \left(s + \frac{m_2}{\sqrt{\alpha}} \pm \sqrt{\alpha} \right)^2 + \frac{2\pi i s m_1}{\sqrt{\alpha}}} ds = \frac{1}{\sqrt{2}} e^{-\frac{\pi \alpha}{2} - \frac{\pi(m_1^2 + m_2^2)}{2\alpha} - \frac{\pi i m_1 m_2}{\alpha} \mp \pi m_2 + \pi i m_1} \end{aligned}$$

and

$$\begin{aligned} \langle f, \pi_{\pm \varepsilon_2} f \rangle_{D^\perp} (m_1 \delta_1 + m_2 \delta_2) &= \int_{\mathbf{R}} \overline{f(s)} f \left(s + \frac{m_2}{\sqrt{\alpha}} \right) e^{\pm 2\pi i \sqrt{\alpha} \left(s + \frac{m_2}{\sqrt{\alpha}} \right) + \frac{2\pi i s m_1}{\sqrt{\alpha}}} ds \\ &= \frac{1}{\sqrt{2}} e^{-\frac{\pi \alpha}{2} - \frac{\pi(m_1^2 + m_2^2)}{2\alpha} - \frac{\pi i m_1 m_2}{\alpha} \mp \pi m_1 + \pi i m_2}, \end{aligned}$$

hence

$$\begin{aligned} \langle f, H_\alpha f \rangle_{D^\perp} &= \frac{1}{\sqrt{2}} \sum_{m_1, m_2} e^{-\frac{\pi \alpha}{2} - \frac{\pi(m_1^2 + m_2^2)}{2\alpha} + \frac{\pi i m_1 m_2}{\alpha}} \\ &\quad \cdot ((e^{\pi m_2} + e^{-\pi m_2})e^{\pi i m_1} + (e^{\pi m_1} + e^{-\pi m_1})e^{\pi i m_2}) V_1^{m_1} V_2^{m_2}. \end{aligned} \tag{3.1}$$

If $\alpha = \frac{1}{q}$, $q \in \mathbf{Z}$, $q \geq 2$, we have $A_{\frac{1}{\alpha}} \simeq C^*(D^{\perp}, \overline{\beta})$ isomorphic to $C(\mathbf{T}^2)$ and $\Theta(eH_{\alpha}e)$ is identified to the function

$$\Theta(eH_{\alpha}e)(z_1, z_2) = \frac{F(z_1, z_2)}{G(z_1, z_2)}, \quad z_1, z_2 \in \mathbf{T},$$

where

$$F(z_1, z_2) = \sum_{m_1, m_2} e^{-\frac{\pi\alpha}{2} - \frac{\pi(m_1^2 + m_2^2)}{2\alpha} + \frac{\pi i m_1 m_2}{\alpha}} \left((e^{\pi m_2} + e^{-\pi m_2}) e^{\pi i m_1} + (e^{\pi m_1} + e^{-\pi m_1}) e^{\pi i m_2} \right) z_1^{m_1} z_2^{m_2},$$

$$G(z_1, z_2) = \sum_{m_1, m_2} e^{-\frac{\pi(m_1^2 + m_2^2)}{2\alpha} + \frac{\pi i m_1 m_2}{\alpha}} z_1^{m_1} z_2^{m_2}.$$

Since

$$\|H_{\alpha}\| \geq \|eH_{\alpha}e\| = \|\Theta(eH_{\alpha}e)\|_{\infty} \geq \frac{F(1, 1)}{G(1, 1)},$$

we further get when q is even

$$\|H_{\frac{1}{q}}\| \geq \phi_0\left(\frac{1}{q}\right) = 4e^{-\frac{\pi}{2q}} \frac{\vartheta\left(\frac{i}{2}, \frac{iq}{2}\right) \vartheta\left(\frac{1}{2}, \frac{iq}{2}\right)}{\vartheta\left(0, \frac{iq}{2}\right)^2} = 4 \frac{\vartheta\left(\frac{1}{q}, \frac{2i}{q}\right) \vartheta\left(\frac{1}{2}, \frac{iq}{2}\right)}{\vartheta\left(0, \frac{2i}{q}\right) \vartheta\left(0, \frac{iq}{2}\right)} \quad (3.2)$$

and when q is odd

$$\|H_{\frac{1}{q}}\| \geq \phi_1\left(\frac{1}{q}\right) = 4e^{-\frac{\pi}{2q}} \frac{\vartheta\left(\frac{i}{2}, \frac{iq}{2}\right) \vartheta\left(\frac{1}{2}, \frac{iq}{2}\right) - 2\vartheta^{odd}\left(\frac{i}{2}, \frac{iq}{2}\right) \vartheta^{odd}\left(\frac{1}{2}, \frac{iq}{2}\right)}{\vartheta\left(0, \frac{iq}{2}\right)^2 - 2\vartheta^{odd}\left(0, \frac{iq}{2}\right)^2}, \quad (3.3)$$

where we set

$$\vartheta^{odd}(z, \tau) = \sum_{m \text{ odd}} e^{\pi i m^2 \tau + 2\pi i m z} = \vartheta(z, \tau) - \vartheta(2z, 4\tau), \quad z \in \mathbf{C} \quad \tau \in \mathbf{H}.$$

Taking as usual

$$\vartheta_{a,b}(z, \tau) = \sum_m e^{\pi \tau (m+a)^2 + 2\pi i (m+a)(z+b)}, \quad z \in \mathbf{C}, \quad \tau \in \mathbf{H}, \quad a, b \in \mathbf{R}, \quad (3.4)$$

we make use of $\vartheta\left(\frac{i}{2}, it\right) = e^{\frac{\pi}{4t}} \vartheta_{\frac{1}{t}, 0}(0, it)$ and $\vartheta\left(\frac{1}{2}, it\right) = \vartheta_{0, \frac{1}{2}}(0, it)$ to get

$$\phi_0(t) = 4 \frac{\vartheta_{0, \frac{1}{2}}\left(0, \frac{i}{2t}\right) \vartheta_{2t, 0}\left(0, \frac{i}{2t}\right)}{\vartheta\left(0, \frac{i}{2t}\right)^2}.$$

The graphs of ϕ_0 and ϕ_1 are drawn below using Mathematica.

```
Plot[4*EllipticTheta[3, Pi*x, Exp[-2*Pi*x]]* EllipticTheta[3, Pi*0.5, Exp[-Pi/(2*x)]]/
(EllipticTheta[3, Pi*0, Exp[-2*Pi*x]]* EllipticTheta[3, Pi*0, Exp[-Pi/(2*x)]]),
{x, 0.01, 0.5}, PlotRange -> {2.7, 4}]
```

Fig. 3.1 The graph of ϕ_0 on $\left(0, \frac{1}{2}\right]$

```

Plot[4*Exp[-Pi*x/2]* (EllipticTheta[3,Pi*0.5,Exp[-Pi/(2*x)]]
*EllipticTheta[3,Pi*I/2,Exp[-Pi/(2*x)]] -2*(EllipticTheta[3,Pi*0.5,Exp[-Pi/(2*x)]]
-EllipticTheta[3,Pi*1,Exp[-2*Pi/x]]) *(EllipticTheta[3,Pi*I/2,Exp[-Pi/(2*x)]]
-EllipticTheta[3,Pi*I,Exp[-2*Pi/x]])/ (EllipticTheta[3,Pi*0,Exp[-Pi/(2*x)]]^2
-2*(EllipticTheta[3,Pi*0,Exp[-Pi/(2*x)]] - EllipticTheta[3,Pi*0,Exp[-2*Pi/x]]^2),
{x,0.01,1/2}, PlotRange -> {2.7,4}]

```

Fig. 3.2 The graph of ϕ_1 on $\left(0, \frac{1}{3}\right]$

They should be compared with Hofstadter's butterfly ([8]).

Estimates (3.2) and (3.3) are quite accurate, as suggested by the following table. The norm of $H_{\frac{1}{q}}$ has been computed using [1, cor.3.2].

α	$\ H_\alpha\ $	$\phi_0(\alpha)$	$\phi_1(\alpha)$
1/2	2.82842	2.82842	
1/3	2.73205		2.73205
1/4	2.82842	2.78648	
1/5	2.96645		2.94109
1/6	3.09557	3.08292	
1/7	3.20330		3.19690
1/8	3.29066	3.28709	
1/9	3.36165		3.35943
1/10	3.42005	3.41855	
1/11	3.46880		3.46771
1/12	3.51004	3.50922	
1/13	3.54537		3.54473
1/50	3.87630	3.87628	
1/51	3.87869		3.87867
1/100	3.93766	3.93765	
1/101	3.93827		3.93827

We notice that inequality (3.2) is sharp for $q = 2$. To see this, remark first that $H_{\frac{1}{2}}^2 = 4 + H_0$, which yields $\|H_{\frac{1}{2}}\| = 2\sqrt{2}$. Using (3.4) and (1.15), we readily see that

$$\vartheta_{0, \frac{1}{2}}(0, i) = \vartheta\left(\frac{1}{2}, i\right) = \vartheta_{\frac{1}{2}, 0}(0, i).$$

On the other hand, Jacobi's identity ([10, p.23]) yields

$$\vartheta(0, i)^4 = \vartheta_{0, \frac{1}{2}}(0, i)^4 + \vartheta_{\frac{1}{2}, 0}(0, i)^4,$$

hence $\vartheta\left(\frac{1}{2}, i\right) = 2^{-\frac{1}{4}}\vartheta(0, i)$ and we get equality in (3.2) in this case.

Notice also that the numerical computations presented above suggest that the estimate is sharp also for $\frac{1}{3}$, which would produce a striking relation between $\vartheta\left(\frac{1}{2}, \frac{3i}{2}\right)$, $\vartheta\left(\frac{1}{3}, \frac{2i}{3}\right)$, $\vartheta\left(\frac{1}{6}, \frac{i}{6}\right)$ and $\vartheta(0, 6i)$.

Next, we derive explicit formulae for the projection e of trace α constructed in proposition 1.1 when $\alpha = \frac{1}{q}$, $q \in \mathbf{Z}$, $q \geq 2$. Again, notation is as in section 1.

For any unitary operator U and $\tau \in \mathbf{H}$, $b \in \mathbf{R}$, $q \in \mathbf{N}^*$, $a \in \mathbf{Z}$, we define the bounded operators

$$\begin{aligned}\vartheta(U, \tau) &= \sum_m e^{\pi i \tau m^2} U^m \\ \vartheta_{0,b}(U, \tau) &= \sum_m e^{\pi i \tau m^2 + 2\pi i m b} U^m \\ \vartheta_{\frac{a}{q}, b}^{(q)}(U, \tau) &= \sum_m e^{\pi i \tau \left(m + \frac{a}{q}\right)^2 + 2\pi i \left(m + \frac{a}{q}\right) b} U^{qm+a}.\end{aligned}$$

The explicit expression of e is given by the following

PROPOSITION 3.1 (i) *If q is even, then*

$$e = \frac{1}{q} \sum_{r,s=0}^{q-1} e^{-\frac{\pi i r s}{q}} \frac{\vartheta_{\frac{s}{q}, \frac{r}{2}}^{(q)}(U_2, \frac{i q}{2}) \vartheta_{\frac{r}{q}, \frac{s}{2}}^{(q)}(U_1, \frac{i q}{2})}{\vartheta(U_2^q, \frac{i q}{2}) \vartheta(U_1^q, \frac{i q}{2})}.$$

(ii) *If q is odd, then*

$$e = \frac{1}{2q} \sum_{r,s=0}^{q-1} e^{-\frac{\pi i r s}{q}} \sum_{\delta_1, \delta_2 \in \{0,1\}} (-1)^{\delta_1 \delta_2} e^{-\frac{\pi i (\delta_2 s + \delta_1 r)}{q}} \frac{\sum_{\varepsilon_1, \varepsilon_2 \in \{0,1\}} (-1)^{\varepsilon_1 \varepsilon_2} \vartheta_{\frac{s}{2q} + \frac{\varepsilon_2}{2}, r + \delta_2}^{(2q)}(U_2, 2iq) \vartheta_{\frac{r}{2q} + \frac{\varepsilon_1}{2}, s + \delta_1}^{(2q)}(U_1, 2iq)}{\sum_{\varepsilon_1, \varepsilon_2 \in \{0,1\}} (-1)^{\varepsilon_1 \varepsilon_2} \vartheta_{\frac{\varepsilon_2}{2}, 0}^{(2)}(U_2, 2iq) \vartheta_{\frac{\varepsilon_1}{2}, 0}^{(2)}(U_1, 2iq)}.$$

Remark. The denominators are central elements in $A_{\frac{1}{q}}$, hence the notation used in the previous formulae makes sense.

Any $b \in C^*(D^\perp, \beta) = A_{\frac{1}{\alpha}}$ is represented as

$$b = \sum_{z \in D^\perp} b(z) \pi_z^* = \sum_{m_1, m_2} b_{m_1, m_2} \pi_{m_1 \delta_1 + m_2 \delta_2}^* = \sum_{m_1, m_2} b_{-m_1, -m_2} e^{\frac{2\pi i m_1 m_2}{\alpha}} V_1^{m_1} V_2^{m_2}. \quad (3.5)$$

For such b and $n_1, n_2 \in \mathbf{Z}$ we set

$$\alpha_{n_1, n_2} = \langle fb, fb \rangle_D (n_1 \sqrt{\alpha}, n_2 \sqrt{\alpha}). \quad (3.6)$$

LEMMA 3.2 *Let $\alpha \in (0, 1)$ and b, α_{n_1, n_2} as in (3.5) and (3.6). Then for all $n_1, n_2 \in \mathbf{Z}$*

$$\begin{aligned}\alpha_{n_1, n_2} &= \frac{1}{\sqrt{2}} e^{-\frac{\pi \alpha (n_1^2 + n_2^2)}{2} + \pi i n_1 n_2 \alpha} \sum_{m_1, \dots, m_4} b_{m_1, m_2} \overline{b_{m_3, m_4}} \\ &\quad \cdot e^{-\frac{\pi ((m_1 - m_3)^2 + (m_2 - m_4)^2)}{2\alpha} - \pi i (n_1 + i n_2) (m_3 - m_1 + i (m_4 - m_2)) + \frac{\pi i (m_1 + m_3)(m_2 - m_4)}{\alpha}}.\end{aligned}$$

Proof. Using

$$(fb)(s) = \sum_{m_1, m_2} b_{m_1, m_2} e^{\frac{2\pi i m_1 m_2}{\alpha} - \pi \left(s - \frac{m_2}{\sqrt{\alpha}}\right)^2 - \frac{2\pi i s m_1}{\sqrt{\alpha}}}, \quad s \in \mathbf{R}$$

and (1.3) we get

$$\begin{aligned}\alpha_{n_1, n_2} &= \int_{\mathbf{R}} (fb)(s) \overline{(fb)(s + n_1 \sqrt{\alpha})} e^{-2\pi i s n_2 \sqrt{\alpha}} ds \\ &= \sum_{m_1, \dots, m_4} b_{m_1, m_2} \overline{b_{m_3, m_4}} e^{\frac{2\pi i (m_1 m_2 - m_3 m_4)}{\alpha} - \frac{\pi (m_2^2 + m_4^2)}{\alpha} - \pi n_1^2 \alpha + 2\pi n_1 m_4} \\ &\quad \cdot \int_{\mathbf{R}} e^{-2\pi s^2 - 2\pi s \left(-\frac{m_2 + m_4}{\sqrt{\alpha}} + n_1 \sqrt{\alpha} + i \left(\frac{m_1 - m_3}{\sqrt{\alpha}} + n_2 \sqrt{\alpha}\right)\right)} ds.\end{aligned}$$

The statement follows now through a plain computation based on (1.16). QED

For all $r, s, n_1, n_2 \in \mathbf{Z}$ we set

$$a_{r,s}^{(n_1, n_2)} = e^{-\frac{\pi(r^2+s^2)}{2\alpha} - \frac{\pi i r s}{\alpha} - \pi i(n_1 + i n_2)(r + i s)} \quad (3.7)$$

$$a^{(n_1, n_2)} = \sum_{r,s} a_{r,s}^{(n_1, n_2)} V_1^r V_2^s. \quad (3.8)$$

Since

$$\sum_{r,s} |a_{r,s}^{(n_1, n_2)}| = \vartheta\left(\frac{i n_1}{2}, \frac{i}{2\alpha}\right) \vartheta\left(\frac{i n_2}{2}, \frac{i}{2\alpha}\right) < \infty,$$

it follows that $a^{(n_1, n_2)} \in C^*(D^\perp, \beta)$. Although we are not going to use it, notice that since $\sigma_{D^\perp}(V_2) = V_1$ and $\sigma_{D^\perp}(V_1) = V_2^{-1}$, we have

$$\sigma_{D^\perp}(a^{(n_1, n_2)}) = a^{(-n_2, n_1)}.$$

If $\alpha = \frac{1}{q}$, $q \in \mathbf{Z}$, $q \geq 2$, lemma 3.2 yields for all $n_1, n_2 \in \mathbf{Z}$

$$\alpha_{n_1, n_2} = \frac{1}{\sqrt{2}} e^{-\frac{\pi(n_1^2 + n_2^2)}{2q} + \frac{\pi i n_1 n_2}{q}} \sum_{m_1, \dots, m_4} b_{m_1, m_2} \overline{b_{m_3, m_4}} a_{m_3 - m_1, m_4 - m_2}^{(n_1, n_2)}. \quad (3.9)$$

Moreover, for any b as in (3.5) we have

$$\begin{aligned} \tau_{D^\perp}(b a^{(n_1, n_2)} b^*) &= \sum_{m_1, \dots, m_4} \sum_{r,s} b_{m_1, m_2} \overline{b_{m_3, m_4}} a_{r,s}^{(n_1, n_2)} \tau_{D^\perp}(V_1^{m_1+r-m_3} V_2^{m_2+s-m_4}) \\ &= \sum_{m_1, \dots, m_4} b_{m_1, m_2} \overline{b_{m_3, m_4}} a_{m_3 - m_1, m_4 - m_2}^{(n_1, n_2)}, \end{aligned}$$

which together with (3.9) gives

$$\begin{aligned} \alpha_{n_1, n_2} &= \frac{1}{\sqrt{2}} e^{-\frac{\pi(n_1^2 + n_2^2)}{2q} + \frac{\pi i n_1 n_2}{q}} \tau_{D^\perp}(b a^{(n_1, n_2)} b^*) \\ &= \frac{1}{\sqrt{2}} e^{-\frac{\pi(n_1^2 + n_2^2)}{2q} + \frac{\pi i n_1 n_2}{q}} \int_{[0,1]^2} a^{(n_1, n_2)}(e^{2\pi i t_1}, e^{2\pi i t_2}) \cdot |b(e^{2\pi i t_1}, e^{2\pi i t_2})|^2 dt_1 dt_2. \end{aligned}$$

In the sequel we will take $b = a^{-\frac{1}{2}} = \langle f, f \rangle_{D^\perp}^{-\frac{1}{2}} \in C(\mathbf{T}^2)$, hence

$$\begin{aligned} \alpha_{n_1, n_2} &= \left\langle f \langle f, f \rangle_{D^\perp}^{-\frac{1}{2}}, f \langle f, f \rangle_{D^\perp}^{-\frac{1}{2}} \right\rangle_D (n_1 \sqrt{\alpha}, n_2 \sqrt{\alpha}) \\ &= \frac{1}{\sqrt{2}} e^{-\frac{\pi(n_1^2 + n_2^2)}{2q} + \frac{\pi i n_1 n_2}{q}} \int_{[0,1]^2} \frac{a^{(n_1, n_2)}(e^{2\pi i t_1}, e^{2\pi i t_2})}{a(e^{2\pi i t_1}, e^{2\pi i t_2})} dt_1 dt_2 \end{aligned} \quad (3.10)$$

and

$$e = |G/D| \sum_{n_1, n_2} \alpha_{n_1, n_2} U_2^{n_2} U_1^{n_1} = \frac{1}{q} \sum_{n_1, n_2} \alpha_{n_1, n_2} U_2^{n_2} U_1^{n_1}. \quad (3.11)$$

For simplicity we shall denote throughout this section

$$\vartheta(z) = \vartheta\left(z, \frac{i q}{2}\right).$$

Proof of proposition 3.1 (i) If q is even, then (1.18) yields for all $t_1, t_2 \in \mathbf{R}$

$$a(e^{2\pi i t_1}, e^{2\pi i t_2}) = \frac{1}{\sqrt{2}} \vartheta(t_1) \vartheta(t_2). \quad (3.12)$$

We employ (3.7) and (3.8) to obtain

$$\alpha^{(n_1, n_2)}(e^{2\pi i t_1}, e^{2\pi i t_2}) = \vartheta\left(t_1 - \frac{n_1 + i n_2}{2}\right) \vartheta\left(t_2 - \frac{n_2 + i n_1}{2}\right). \quad (3.13)$$

Writing $n_1 = ql_1 + r$, $n_2 = ql_2 + s$, $l_1, l_2, r, s \in \mathbf{Z}$, $0 \leq r, s < q$ and employing (3.10), (3.12) and (3.13) we get

$$\alpha_{n_1, n_2} = e^{-\frac{\pi q(l_1^2 + l_2^2)}{2} - \pi(l_1 r + l_2 s) - \frac{\pi(r^2 + s^2)}{2q} + \pi i(l_1 s + l_2 r) + \frac{\pi i r s}{q}} \int_0^1 \frac{\vartheta\left(t_1 - \frac{r + i s + i l_2 q}{2}\right)}{\vartheta(t_1)} dt_1 \int_0^1 \frac{\vartheta\left(t_2 - \frac{s + i r + i l_1 q}{2}\right)}{\vartheta(t_2)} dt_2.$$

Using also the quasiperiodicity of ϑ :

$$\vartheta\left(z + \frac{i l q}{2}\right) = e^{\frac{\pi q l^2}{2} - 2\pi i l z} \vartheta(z) \quad z \in \mathbf{C}, \quad l \in \mathbf{Z},$$

we get

$$\alpha_{ql_1 + r, ql_2 + s} = e^{-\frac{\pi(r^2 + s^2)}{2q} + \frac{\pi i r s}{q}} \int_0^1 \frac{e^{2\pi i l_2 t_1} \vartheta\left(t_1 - \frac{r + i s}{2}\right)}{\vartheta(t_1)} dt_1 \int_0^1 \frac{e^{2\pi i l_1 t_2} \vartheta\left(t_2 - \frac{s + i r}{2}\right)}{\vartheta(t_2)} dt_2.$$

Furthermore, using also

$$\sum_l e^{2\pi i l y} \int_0^1 \frac{e^{2\pi i l x} \vartheta\left(x - \frac{r + i s}{2}\right)}{\vartheta(x)} dx = \sum_l \int_0^1 \frac{e^{2\pi i l x} \vartheta\left(x - y - \frac{r + i s}{2}\right)}{\vartheta(x - y)} dx = \frac{\vartheta\left(y + \frac{r + i s}{2}\right)}{\vartheta(y)}$$

and the fact that U_1^q and U_2^q are central in $A_{\frac{1}{q}}$, we get

$$\sum_{l_1, l_2} \alpha_{ql_1 + r, ql_2 + s} U_2^{ql_2} U_1^{ql_1} = e^{-\frac{\pi(r^2 + s^2)}{2q} + \frac{\pi i r s}{q}} \cdot \frac{\vartheta\left(e^{\pi i(r + i s)} U_2^q, \frac{i q}{2}\right) \vartheta\left(e^{\pi i(s + i r)} U_1^q, \frac{i q}{2}\right)}{\vartheta\left(U_2^q, \frac{i q}{2}\right) \vartheta\left(U_1^q, \frac{i q}{2}\right)}.$$

The equality from proposition 3.1 (i) follows from this and from

$$e^{-\frac{\pi s^2}{2q} + \frac{\pi i r s}{q}} U^s \vartheta\left(e^{\pi i(r + i s)} U^q, \frac{i q}{2}\right) = \vartheta_{\frac{s}{q}, \frac{r}{2}}^{(q)}\left(U, \frac{i q}{2}\right). \quad \text{QED} \quad (3.14)$$

Before completing the proof of proposition 3.1 for q odd, we introduce some notation by setting for all $z \in \mathbf{C}$, $\tau \in \mathbf{H}$, $a, b \in \mathbf{R}$

$$\vartheta_{a,b}^{odd}(z, \tau) = \sum_{m \text{ odd}} e^{\pi i \tau(m+a)^2 + 2\pi i(m+a)(z+b)} = \vartheta_{a,b}(z, \tau) - \vartheta_{\frac{a}{2}, 2b}(2z, 4\tau) = \vartheta_{\frac{a+1}{2}, 2b}(2z, 4\tau),$$

$$\vartheta_{a,b}^{even} = \sum_{m \text{ even}} e^{\pi i \tau(m+a)^2 + 2\pi i(m+a)(z+b)} = \vartheta_{\frac{a}{2}, 2b}(2z, 4\tau),$$

$$\vartheta^\diamond(z, \tau) = \vartheta_{0,0}^\diamond(z, \tau), \quad \diamond \in \{\text{even}, \text{odd}\}.$$

For all $z \in \mathbf{C}$, $\tau \in \mathbf{H}$, $a, b \in \mathbf{R}$, $l \in \mathbf{Z}$ we have

$$\vartheta^{even}\left(z + \frac{1}{2}, \tau\right) = \vartheta^{even}(z, \tau), \quad (3.15)$$

$$\vartheta^{odd}\left(z + \frac{1}{2}, \tau\right) = -\vartheta^{odd}(z, \tau), \quad (3.16)$$

$$\vartheta_{a,b}^{odd}(z + l\tau, \tau) = e^{-\pi i \tau l^2 - 2\pi i l(z+b)} \cdot \begin{cases} \vartheta_{a,b}^{odd}(z, \tau), & \text{if } l \text{ is even} \\ \vartheta_{a,b}^{even}(z, \tau), & \text{if } l \text{ is odd} \end{cases}. \quad (3.17)$$

The following lemma is easy to prove.

LEMMA 3.3 *Let $\phi : \mathbf{R}^2 \rightarrow \mathbf{C}$ be a continuous function which is periodic modulo $\mathbf{Z}^2$. Then*

$$\begin{aligned} \sum_{l_1, l_2 \text{ even}} \int_{[0,1]^2} e^{2\pi i(l_1 t_1 + l_2 t_2)} \phi(t_1, t_2) dt_1 dt_2 &= \frac{1}{4} \left(\phi(0, 0) + \phi\left(\frac{1}{2}, 0\right) + \phi\left(0, \frac{1}{2}\right) + \phi\left(\frac{1}{2}, \frac{1}{2}\right) \right), \\ \sum_{l_1 \text{ even}, l_2 \text{ odd}} \int_{[0,1]^2} e^{2\pi i(l_1 t_1 + l_2 t_2)} \phi(t_1, t_2) dt_1 dt_2 &= \frac{1}{4} \left(\phi(0, 0) + \phi\left(\frac{1}{2}, 0\right) - \phi\left(0, \frac{1}{2}\right) - \phi\left(\frac{1}{2}, \frac{1}{2}\right) \right), \\ \sum_{l_1 \text{ odd}, l_2 \text{ even}} \int_{[0,1]^2} e^{2\pi i(l_1 t_1 + l_2 t_2)} \phi(t_1, t_2) dt_1 dt_2 &= \frac{1}{4} \left(\phi(0, 0) - \phi\left(\frac{1}{2}, 0\right) + \phi\left(0, \frac{1}{2}\right) - \phi\left(\frac{1}{2}, \frac{1}{2}\right) \right), \\ \sum_{l_1, l_2 \text{ odd}} \int_{[0,1]^2} e^{2\pi i(l_1 t_1 + l_2 t_2)} \phi(t_1, t_2) dt_1 dt_2 &= \frac{1}{4} \left(\phi(0, 0) - \phi\left(\frac{1}{2}, 0\right) - \phi\left(0, \frac{1}{2}\right) + \phi\left(\frac{1}{2}, \frac{1}{2}\right) \right). \end{aligned}$$

Proof. We consider for instance the second equality. The others are similar. if we set $l_1 = 2m_1$, $l_2 = 2m_2 + 1$, $x_1 = 2t_1$, $x_2 = 2t_2$, and divide $[0, 2]^2$ into four equal squares, then we get :

$$\begin{aligned} S_{01} &= \sum_{l_1 \text{ even}, l_2 \text{ odd}} \int_{[0,1]^2} e^{2\pi i(l_1 t_1 + l_2 t_2)} \phi(t_1, t_2) dt_1 dt_2 \\ &= \frac{1}{4} \sum_{m_1, m_2} \int_{[0,2]^2} e^{2\pi i(m_1 x_1 + m_2 x_2)} e^{\pi i x_2} \phi\left(\frac{x_1}{2}, \frac{x_2}{2}\right) dx_1 dx_2 \\ &= \frac{1}{4} \sum_{m_1, m_2} \int_{[0,1]^2} e^{2\pi i(m_1 x_1 + m_2 x_2)} \psi(x_1, x_2) dx_1 dx_2, \end{aligned}$$

where

$$\psi(x_1, x_2) = e^{\pi i x_2} \phi\left(\frac{x_1}{2}, \frac{x_2}{2}\right) + e^{\pi i x_2} \phi\left(\frac{x_1+1}{2}, \frac{x_2}{2}\right) - e^{\pi i x_2} \phi\left(\frac{x_1}{2}, \frac{x_2+1}{2}\right) - e^{\pi i x_2} \phi\left(\frac{x_1+1}{2}, \frac{x_2+1}{2}\right).$$

Since $\psi(x_1 + 1, x_2) = \psi(x_1, x_2) = \psi(x_1, x_2 + 1)$, $x_1, x_2 \in \mathbf{R}$, we get

$$S_{01} = \frac{1}{4} \psi(0, 0) = \frac{1}{4} \left(\phi(0, 0) + \phi\left(\frac{1}{2}, 0\right) - \phi\left(0, \frac{1}{2}\right) - \phi\left(\frac{1}{2}, \frac{1}{2}\right) \right). \quad \text{QED}$$

For all $a, b \in \mathbf{R}$, $z \in \mathbf{C}$, we set

$$\vartheta_{a,b}(z) = \vartheta_{a,b}\left(z, \frac{iq}{2}\right), \quad \vartheta_{a,b}^\diamond(z) = \vartheta_{a,b}^\diamond\left(z, \frac{iq}{2}\right), \quad \diamond \in \{\text{even}, \text{odd}\},$$

and consider the following continuous functions on $\mathbf{R}^2/\mathbf{Z}^2$

$$\begin{aligned} \phi_{r,s}^{0,0}(t_1, t_2) &= \frac{\vartheta\left(t_1 + \frac{r+is}{2}\right) \vartheta\left(t_2 + \frac{s+ir}{2}\right) - 2 \vartheta^{\text{odd}}\left(t_1 + \frac{r+is}{2}\right) \vartheta^{\text{odd}}\left(t_2 + \frac{s+ir}{2}\right)}{\vartheta(t_1) \vartheta(t_2) - 2 \vartheta^{\text{odd}}(t_1) \vartheta^{\text{odd}}(t_2)}, \\ \phi_{r,s}^{0,1}(t_1, t_2) &= \frac{\vartheta\left(t_1 + \frac{r+is}{2}\right) \vartheta_{0,\frac{1}{2}}\left(t_2 + \frac{s+ir}{2}\right) + 2 \vartheta^{\text{even}}\left(t_1 + \frac{r+is}{2}\right) \vartheta^{\text{odd}}\left(t_2 + \frac{s+ir}{2}\right)}{\vartheta(t_1) \vartheta(t_2) - 2 \vartheta^{\text{odd}}(t_1) \vartheta^{\text{odd}}(t_2)}, \\ \phi_{r,s}^{1,0}(t_1, t_2) &= \frac{\vartheta_{0,\frac{1}{2}}\left(t_1 + \frac{r+is}{2}\right) \vartheta\left(t_2 + \frac{s+ir}{2}\right) + 2 \vartheta^{\text{odd}}\left(t_1 + \frac{r+is}{2}\right) \vartheta^{\text{even}}\left(t_2 + \frac{s+ir}{2}\right)}{\vartheta(t_1) \vartheta(t_2) - 2 \vartheta^{\text{odd}}(t_1) \vartheta^{\text{odd}}(t_2)}, \\ \phi_{r,s}^{1,1}(t_1, t_2) &= \frac{\vartheta_{0,\frac{1}{2}}\left(t_1 + \frac{r+is}{2}\right) \vartheta_{0,\frac{1}{2}}\left(t_2 + \frac{s+ir}{2}\right) - 2 \vartheta^{\text{even}}\left(t_1 + \frac{r+is}{2}\right) \vartheta^{\text{even}}\left(t_2 + \frac{s+ir}{2}\right)}{\vartheta(t_1) \vartheta(t_2) - 2 \vartheta^{\text{odd}}(t_1) \vartheta^{\text{odd}}(t_2)}. \end{aligned}$$

Proof of proposition 3.1 (ii). Our first aim is to compute $\sum_{l_1, l_2} \alpha_{ql_1+r, ql_2+s} U_2^{ql_2} U_1^{ql_1}$. This is split into four sums as follows :

$$\begin{aligned} S_{00} &= \sum_{l_1, l_2 \text{ even}} \alpha_{ql_1+r, ql_2+s} e^{2\pi i(l_1 x_1 + l_2 x_2)}, & S_{01} &= \sum_{l_1 \text{ even}, l_2 \text{ odd}} \alpha_{ql_1+r, ql_2+s} e^{2\pi i(l_1 x_1 + l_2 x_2)}, \\ S_{10} &= \sum_{l_1 \text{ odd}, l_2 \text{ even}} \alpha_{ql_1+r, ql_2+s} e^{2\pi i(l_1 x_1 + l_2 x_2)}, & S_{11} &= \sum_{l_1, l_2 \text{ odd}} \alpha_{ql_1+r, ql_2+s} e^{2\pi i(l_1 x_1 + l_2 x_2)}. \end{aligned}$$

From (1.18) we get for all $t_1, t_2 \in \mathbf{R}$

$$a(e^{2\pi i t_1}, e^{2\pi i t_2}) = \frac{1}{\sqrt{2}} (\vartheta(t_1) \vartheta(t_2) - 2 \vartheta^{\text{odd}}(t_1) \vartheta^{\text{odd}}(t_2)) > 0, \quad (3.18)$$

whilst from (3.7) and (3.8)

$$\begin{aligned} a^{(n_1, n_2)}(e^{2\pi i t_1}, e^{2\pi i t_2}) &= \frac{1}{\sqrt{2}} \left(\vartheta\left(t_1 - \frac{n_1 + i n_2}{2}\right) \vartheta\left(t_2 - \frac{n_2 + i n_1}{2}\right) \right. \\ &\quad \left. - 2 \vartheta^{\text{odd}}\left(t_1 - \frac{n_1 + i n_2}{2}\right) \vartheta^{\text{odd}}\left(t_2 - \frac{n_2 + i n_1}{2}\right) \right). \end{aligned} \quad (3.19)$$

To compute S_{00} , we notice that the quasi-periodicity properties (3.15), (3.16), (3.17) and

$$\vartheta\left(z + \frac{ilq}{2}\right) = e^{\frac{\pi q l^2}{2} - 2\pi i l z} \vartheta(z), \quad z \in \mathbf{C}, \quad l \in \mathbf{Z} \quad (3.20)$$

yield for all $t_1, t_2 \in \mathbf{R}$, $n_1, n_2 \in \mathbf{Z}$, $n_1 = ql_1 + r$, $n_2 = ql_2 + s$, $l_1, l_2, r, s \in \mathbf{Z}$, $0 \leq r, s < q$

$$\begin{aligned} a^{(n_1, n_2)}(e^{2\pi i t_1}, e^{2\pi i t_2}) &= \frac{1}{\sqrt{2}} e^{\frac{\pi(l_1^2 + l_2^2)q}{2} - 2\pi i(l_2 t_1 + l_1 t_2) - \pi i(l_2 r + l_1 s) + \pi(l_2 s + l_1 r)} \\ &\quad \cdot \left(\vartheta\left(t_1 - \frac{r + is}{2}\right) \vartheta\left(t_2 - \frac{s + ir}{2}\right) - 2 \vartheta^{\text{odd}}\left(t_1 - \frac{r + is}{2}\right) \vartheta^{\text{odd}}\left(t_2 - \frac{s + ir}{2}\right) \right). \end{aligned} \quad (3.21)$$

We combine (3.10), (3.18), (3.21) and the definition of $\phi_{r,s}^{0,0}$ to get

$$\alpha_{n_1, n_2} = e^{-\frac{\pi(r^2 + s^2)}{2q} + \frac{\pi i r s}{q}} \int_{[0,1]^2} e^{2\pi i(l_2 t_1 + l_1 t_2)} \phi_{r,s}^{0,0}(-t_1, -t_2) dt_1 dt_2.$$

Using lemma 3.3 we gather

$$\begin{aligned} S_{00} &= e^{-\frac{\pi(r^2 + s^2)}{2q} + \frac{\pi i r s}{q}} \sum_{l_1, l_2 \text{ even}} \int_{[0,1]^2} e^{2\pi i l_2(t_1 + x_2) + 2\pi i l_1(t_2 + x_1)} \phi_{r,s}^{0,0}(-t_1, -t_2) dt_1 dt_2 \\ &= e^{-\frac{\pi(r^2 + s^2)}{2q} + \frac{\pi i r s}{q}} \sum_{l_1, l_2 \text{ even}} \int_{[0,1]^2} e^{2\pi i(l_1 t_1 + l_2 t_2)} \phi_{r,s}^{0,0}(x_2 - t_2, x_1 - t_1) dt_1 dt_2 \\ &= \frac{1}{4} e^{-\frac{\pi(r^2 + s^2)}{2q} + \frac{\pi i r s}{q}} \left(\phi_{r,s}^{0,0}(x_2, x_1) + \phi_{r,s}^{0,0}\left(x_2, x_1 - \frac{1}{2}\right) + \phi_{r,s}^{0,0}\left(x_2 - \frac{1}{2}, x_1\right) \right. \\ &\quad \left. + \phi_{r,s}^{0,0}\left(x_2 - \frac{1}{2}, x_1 - \frac{1}{2}\right) \right). \end{aligned} \quad (3.22)$$

Similar computations yield

$$S_{01} = \frac{1}{4} e^{-\frac{\pi(r^2+s^2)}{2q} + \frac{\pi i r s}{q}} \left(\phi_{r,s}^{0,1}(x_2, x_1) + \phi_{r,s}^{0,1}\left(x_2, x_1 - \frac{1}{2}\right) - \phi_{r,s}^{0,1}\left(x_2 - \frac{1}{2}, x_1\right) - \phi_{r,s}^{0,1}\left(x_2 - \frac{1}{2}, x_1 - \frac{1}{2}\right) \right). \quad (3.23)$$

$$S_{10} = \frac{1}{4} e^{-\frac{\pi(r^2+s^2)}{2q} + \frac{\pi i r s}{q}} \left(\phi_{r,s}^{1,0}(x_2, x_1) - \phi_{r,s}^{1,0}\left(x_2, x_1 - \frac{1}{2}\right) + \phi_{r,s}^{1,0}\left(x_2 - \frac{1}{2}, x_1\right) - \phi_{r,s}^{1,0}\left(x_2 - \frac{1}{2}, x_1 - \frac{1}{2}\right) \right). \quad (3.24)$$

$$S_{11} = \frac{1}{4} e^{-\frac{\pi(r^2+s^2)}{2q} + \frac{\pi i r s}{q}} \left(\phi_{r,s}^{1,1}(x_2, x_1) - \phi_{r,s}^{1,1}\left(x_2, x_1 - \frac{1}{2}\right) - \phi_{r,s}^{1,1}\left(x_2 - \frac{1}{2}, x_1\right) + \phi_{r,s}^{1,1}\left(x_2 - \frac{1}{2}, x_1 - \frac{1}{2}\right) \right). \quad (3.25)$$

Immediate computations based on the obvious equalities

$$\vartheta(z, \tau) = \vartheta^{even}(z, \tau) + \vartheta^{odd}(z, \tau) \quad \text{and} \quad \vartheta_{0, \frac{1}{2}}(z, \tau) = \vartheta^{even}(z, \tau) - \vartheta^{odd}(z, \tau)$$

show that $\phi_{r,s}^{0,0} = \phi_{r,s}^{0,1} = \phi_{r,s}^{1,0} = -\phi_{r,s}^{1,1} = \phi_{r,s}$ and actually we get for all $r, s \in \mathbf{Z}$

$$\phi_{r,s}(t_1, t_2) = \frac{\vartheta\left(t_1 + \frac{r+is}{2}\right) \vartheta\left(t_2 + \frac{s+ir}{2}\right) - 2 \vartheta^{odd}\left(t_1 + \frac{r+is}{2}\right) \vartheta^{odd}\left(t_2 + \frac{s+ir}{2}\right)}{\vartheta(t_1) \vartheta(t_2) - 2 \vartheta^{odd}(t_1) \vartheta^{odd}(t_2)}. \quad (3.26)$$

From (3.22)-(3.25) we get

$$\begin{aligned} \sum_{l_1, l_2} \alpha_{ql_1+r, ql_2+s} e^{2\pi i(l_1 x_1 + l_2 x_2)} &= S_{00} + S_{01} + S_{10} + S_{11} \\ &= \frac{1}{2} e^{-\frac{\pi(r^2+s^2)}{2q} + \frac{\pi i r s}{q}} \sum_{\delta_1, \delta_2 \in \{0,1\}} (-1)^{\delta_1 \delta_2} \phi_{r,s}\left(x_2 - \frac{\delta_2}{2}, x_1 - \frac{\delta_1}{2}\right). \end{aligned} \quad (3.27)$$

We identify U_1^q with $e^{2\pi i x_1}$ and U_2^q with $e^{2\pi i x_2}$ to obtain

$$\begin{aligned} \sum_{l_1, l_2} \alpha_{ql_1+r, ql_2+s} U_2^{ql_2} U_1^{ql_1} &= \frac{1}{2} e^{-\frac{\pi(r^2+s^2)}{2q} + \frac{\pi i r s}{q}} \left(\vartheta\left(U_2^q, \frac{iq}{2}\right) \vartheta\left(U_1^q, \frac{iq}{2}\right) - 2 \vartheta^{odd}\left(U_2^q, \frac{iq}{2}\right) \vartheta^{odd}\left(U_1^q, \frac{iq}{2}\right) \right)^{-1} \\ &\cdot \sum_{\delta_1, \delta_2 \in \{0,1\}} (-1)^{\delta_1 \delta_2} \left(\vartheta\left(e^{\pi i(r+\delta_2+is)} U_2^q, \frac{iq}{2}\right) \vartheta\left(e^{\pi i(s+\delta_1+ir)} U_1^q, \frac{iq}{2}\right) \right. \\ &\quad \left. - 2 \vartheta^{odd}\left(e^{\pi i(r+\delta_2+is)} U_2^q, \frac{iq}{2}\right) \vartheta^{odd}\left(e^{\pi i(s+\delta_1+ir)} U_1^q, \frac{iq}{2}\right) \right), \end{aligned} \quad (3.28)$$

where we denote for any unitary U and $\tau \in \mathbf{H}$, $q \in \mathbf{N}^*$, $a \in \mathbf{Z}$, $b \in \mathbf{R}$

$$\vartheta^{odd}(z, \tau) = \sum_{m \text{ odd}} e^{\pi i \tau m^2} U^m, \quad \vartheta^{even}(z, \tau) = \sum_{m \text{ even}} e^{\pi i \tau m^2} U^m$$

and

$$\vartheta_{\frac{a}{q},b}^{(q) odd}(U, \tau) = \sum_{m \text{ odd}} e^{\pi i \tau \left(m + \frac{a}{q}\right)^2 + 2\pi i \left(m + \frac{a}{q}\right)b} U^{qm+a} = \vartheta_{\frac{a}{2q} + \frac{1}{2}, 2b}^{(2q)}(U, 4\tau), \quad (3.29)$$

$$\vartheta_{\frac{a}{q},b}^{(q) even}(U, \tau) = \sum_{m \text{ even}} e^{\pi i \tau \left(m + \frac{a}{q}\right)^2 + 2\pi i \left(m + \frac{a}{q}\right)b} U^{qm+a} = \vartheta_{\frac{a}{2q}, 2b}^{(2q)}(U, 4\tau). \quad (3.30)$$

Using $\vartheta(U, \tau) = \vartheta^{odd}(U, \tau) + \vartheta^{even}(U, \tau)$, $\vartheta^{odd}(U, \tau) = \vartheta_{\frac{1}{2}, 0}^{(2)}(U, 4\tau)$ and $\vartheta^{even}(U, \tau) = \vartheta_{0, 0}^{(2)}(U, 4\tau)$, we get

$$\begin{aligned} & \vartheta\left(U_2^q, \frac{iq}{2}\right) \vartheta\left(U_1^q, \frac{iq}{2}\right) - 2\vartheta^{odd}\left(U_2^q, \frac{iq}{2}\right) \vartheta^{odd}\left(U_1^q, \frac{iq}{2}\right) \\ &= \sum_{\varepsilon_1, \varepsilon_2 \in \{0, 1\}} (-1)^{\varepsilon_1 \varepsilon_2} \vartheta_{\frac{\varepsilon_2}{2}, 0}^{(2)}(U_2^q, 2iq) \vartheta_{\frac{\varepsilon_1}{2}, 0}^{(2)}(U_1^q, 2iq). \end{aligned} \quad (3.31)$$

Using the quasi-periodicity relations (3.14) and

$$e^{-\frac{\pi s^2}{2q} + \frac{\pi i r s}{q}} U^s \vartheta^{odd}\left(e^{\pi i(r+is)} U^q, \frac{iq}{2}\right) = \vartheta_{\frac{s}{q}, \frac{r}{2}}^{(q) odd}\left(U, \frac{iq}{2}\right),$$

and also $\vartheta_{\frac{a}{q},b}^{(q)}(U, \tau) = \vartheta_{\frac{a}{q},b}^{(q) even}(U, \tau) + \vartheta_{\frac{a}{q},b}^{(q) odd}(U, \tau)$, (3.29) and (3.30), we get

$$\begin{aligned} & e^{-\frac{\pi(r^2+s^2)}{2q} + \frac{\pi i r s}{q}} U_2^s \left(\vartheta\left(e^{\pi i(r+\delta_2+is)} U_2^q, \frac{iq}{2}\right) \vartheta\left(e^{\pi i(s+\delta_1+ir)} U_1^q, \frac{iq}{2}\right) \right. \\ & \quad \left. - 2\vartheta^{odd}\left(e^{\pi i(r+\delta_2+is)} U_2^q, \frac{iq}{2}\right) \vartheta^{odd}\left(e^{\pi i(s+\delta_1+ir)} U_1^q, \frac{iq}{2}\right) \right) U_1^r \\ &= e^{-\frac{\pi i r s}{q} - \frac{\pi i(\delta_2 s + \delta_1 r)}{q}} \sum_{\varepsilon_1, \varepsilon_2 \in \{0, 1\}} (-1)^{\varepsilon_1 \varepsilon_2} \vartheta_{\frac{s}{2q} + \frac{\varepsilon_2}{2}, r+\delta_2}^{(2q)}(U_2, 2iq) \vartheta_{\frac{r}{2q} + \frac{\varepsilon_1}{2}, s+\delta_1}^{(2q)}(U_1, 2iq). \end{aligned} \quad (3.32)$$

Proposition 3.1 (ii) follows now from (3.11), (3.28), (3.31) and (3.32). QED

We continue with some considerations on the case $\alpha = \frac{1}{2}$. According to proposition 3.1, the formula

$$e = \frac{1}{2} \left(1 + \frac{\vartheta_{0, \frac{1}{2}}^{(2)}(U_2, i) \vartheta_{\frac{1}{2}, 0}^{(2)}(U_1, i)}{\vartheta(U_2^2, i) \vartheta(U_1^2, i)} + \frac{\vartheta_{\frac{1}{2}, 0}^{(2)}(U_2, i) \vartheta_{0, \frac{1}{2}}^{(2)}(U_1, i)}{\vartheta(U_2^2, i) \vartheta(U_1^2, i)} - \frac{i \vartheta_{\frac{1}{2}, \frac{1}{2}}^{(2)}(U_2, i) \vartheta_{\frac{1}{2}, \frac{1}{2}}^{(2)}(U_1, i)}{\vartheta(U_2^2, i) \vartheta(U_1^2, i)} \right)$$

defines a projection of trace $\frac{1}{2}$ in $A_{\frac{1}{2}}$. We shall simply denote $e = e_1 = (+ \ + \ + \ -)$.

We consider the automorphisms ρ_{t_1, t_2} of $A_{\frac{1}{2}}$ acting on the generators U_1 and U_2 by $\rho_{t_1, t_2}(U_j) = e^{2\pi i t_j}$, $j = 1, 2$, $t_1, t_2 \in \mathbf{R}$ and ρ acting by $\rho(U_1) = -U_2$, $\rho(U_2) = -U_1$. The relations

$$\vartheta_{\frac{1}{2}, b}^{(2)}(-U, \tau) = -\vartheta_{\frac{1}{2}, b}^{(2)}(U, \tau), \quad b \in \mathbf{R}, \tau \in \mathbf{H},$$

$$\vartheta_{\frac{1}{2}, \frac{1}{2}}^{(2)}(U_1, \tau) \vartheta_{\frac{1}{2}, \frac{1}{2}}^{(2)}(U_2, \tau) = -\vartheta_{\frac{1}{2}, \frac{1}{2}}^{(2)}(U_2, \tau) \vartheta_{\frac{1}{2}, \frac{1}{2}}^{(2)}(U_1, \tau),$$

and the fact that $\vartheta(U_1^2, i)$ and $\vartheta(U_2^2, i)$ are central in $A_{\frac{1}{2}}$ show that

$$e_1 = (+ \ + \ + \ -),$$

$$e_2 = \rho_{\frac{1}{2}, 0}(e_1) = (+ \ - \ + \ +),$$

$$e_3 = \rho_{0, \frac{1}{2}}(e_1) = (+ \ + \ - \ +),$$

$$e_4 = \rho_{\frac{1}{2}, \frac{1}{2}}(e_1) = (+ \ - \ - \ -),$$

$$e_5 = \rho(e_1) = (+ \ - \ - \ +).$$

In particular e_j , $1 \leq j \leq 5$ are projections of trace $\frac{1}{2}$ in $A_{\frac{1}{2}}$ such that

$$e_1 + e_5 = e + \rho(e) = 1, \quad (3.33)$$

$$e_1 + e_2 + e_3 + e_4 = \sum_{c_1, c_2=0}^1 \rho_{\frac{c_1}{2}, \frac{c_2}{2}}(e) = 2. \quad (3.34)$$

We also notice that if $\tilde{\rho}$ is the automorphism of $A_{\frac{1}{2}}$ defined by $\tilde{\rho}(U_1) = U_2$, $\tilde{\rho}(U_2) = U_1$, then

$$\tilde{\rho}(e_1) = (+ \ + \ + \ +).$$

Actually, the analog of (3.34) holds for any $\alpha = \frac{1}{q}$, $q \in \mathbf{N}^*$, $q \geq 2$. To see this, we notice first that for all $s, c \in \mathbf{Z}$, $b \in \mathbf{R}$

$$\vartheta_{\frac{s}{q}, b}^{(q)}(e^{\frac{2\pi i c}{q}} U, \tau) = e^{\frac{2\pi i s c}{q}} \vartheta_{\frac{s}{q}, b}^{(q)}(U, \tau),$$

This implies that if we write $e = \sum_{r,s=0}^{q-1} e^{-\frac{\pi i r s}{q}} A_{r,s}$ as in proposition 3.1, then

$$\rho_{\frac{c_1}{q}, \frac{c_2}{q}}(e) = \frac{1}{q} \sum_{r,s=0}^{q-1} e^{-\frac{\pi i r s}{q} + \frac{2\pi i s c_2}{q} + \frac{2\pi i r c_1}{q}} A_{r,s}$$

and furthermore

$$\sum_{c_1, c_2=0}^{q-1} \rho_{\frac{c_1}{q}, \frac{c_2}{q}}(e) = \frac{1}{q} \sum_{r,s=0}^{q-1} e^{-\frac{\pi i r s}{q}} \left(\sum_{c_1=0}^{q-1} e^{\frac{2\pi i r c_1}{q}} \right) \left(\sum_{c_2=0}^{q-1} e^{\frac{2\pi i s c_2}{q}} \right) A_{r,s} = q A_{0,0} = q. \quad (3.35)$$

Let $\pi : A_{\frac{1}{q}} \rightarrow M_q(\mathbf{C})$, $\pi(U_j) = \tilde{U}_j$, $j = 1, 2$ be the canonical finite dimensional representation of $A_{\frac{1}{q}}$ (i.e. $\tilde{U}_1^q = \tilde{U}_2^q = I_q$). We consider the finite dimensional representations $\pi_{t_1, t_2} = \pi \rho_{t_1, t_2}$ of $A_{\frac{1}{q}}$. Then, we get $\pi_{t_1, t_2}(U_j^{qm+s}) = e^{2\pi i(qm+s)t_j} \tilde{U}_j^s$, $j = 1, 2$, $0 \leq s < q$ and

$$\pi_{t_1, t_2} \left(\vartheta_{\frac{s}{q}, \frac{r}{2}}^{(q)}(U_2, \tau) \right) = \sum_m e^{\pi i \tau \left(m + \frac{s}{q}\right)^2 + 2\pi i \left(m + \frac{s}{q}\right) \frac{r}{2}} e^{2\pi i(qm+s)t_2} \tilde{U}_2^s = \vartheta_{\frac{s}{q}, \frac{r}{2}}(qt_2, \tau) \tilde{U}_2^s,$$

$$\pi_{t_1, t_2} \left(\vartheta_{\frac{r}{q}, \frac{s}{2}}^{(q)}(U_1, \tau) \right) = \vartheta_{\frac{r}{q}, \frac{s}{2}}(qt_1, \tau) \tilde{U}_1^r$$

$$\pi_{t_1, t_2}(\vartheta(U_j^q, \tau)) = \vartheta(qt_j, \tau) \cdot I_q, \quad j = 1, 2.$$

Therefore, for any non-negative even integer q

$$\pi_{t_1, t_2}(e) = \frac{\sum_{r,s=0}^{q-1} e^{-\frac{\pi i r s}{q}} \vartheta_{\frac{s}{q}, \frac{r}{2}}(qt_2, \frac{iq}{2}) \vartheta_{\frac{r}{q}, \frac{s}{2}}(qt_1, \frac{iq}{2}) \tilde{U}_2^s \tilde{U}_1^r}{q \vartheta(qt_2, \frac{iq}{2}) \vartheta(qt_1, \frac{iq}{2})}. \quad (3.36)$$

Using (3.36) we compute $\tau(\pi_{t_1, t_2}(e)^2)$. As this is equal to $\tau(\pi_{t_1, t_2}(e)) = q^{-1}$, we get for all $t_1, t_2 \in \mathbf{R}$

$$\begin{aligned} & \sum_{m,n=0}^{q-1} e^{-\frac{4\pi i m n}{q}} \vartheta_{-\frac{n}{q}, -\frac{m}{2}} \left(t_1, \frac{iq}{2} \right) \vartheta_{\frac{n}{q}, \frac{m}{2}} \left(t_1, \frac{iq}{2} \right) \vartheta_{-\frac{m}{q}, -\frac{n}{2}} \left(t_2, \frac{iq}{2} \right) \vartheta_{\frac{m}{q}, \frac{n}{2}} \left(t_2, \frac{iq}{2} \right) \\ &= q \vartheta \left(t_1, \frac{iq}{2} \right)^2 \vartheta \left(t_2, \frac{iq}{2} \right)^2. \end{aligned} \quad (3.37)$$

Using $\vartheta_{-a,-b}(z, \tau) = \vartheta_{a,b}(-z, \tau)$, it follows that for any even integer $q \in \mathbf{N}^*$ and any $t_1, t_2 \in \mathbf{R}$

$$\begin{aligned} \sum_{m,n=0}^{q-1} e^{-\frac{4\pi i m n}{q}} \vartheta_{\frac{n}{q}, \frac{m}{2}}\left(t_1, \frac{i q}{2}\right) \vartheta_{\frac{n}{q}, \frac{m}{2}}\left(-t_1, \frac{i q}{2}\right) \vartheta_{\frac{m}{q}, \frac{n}{2}}\left(t_2, \frac{i q}{2}\right) \vartheta_{\frac{m}{q}, \frac{n}{2}}\left(-t_2, \frac{i q}{2}\right) \\ = q \vartheta\left(t_1, \frac{i q}{2}\right)^2 \vartheta\left(t_2, \frac{i q}{2}\right)^2. \end{aligned} \quad (3.38)$$

Actually (3.38) can be regarded as a sort of Riemann theta relation. To see this, we specialize further to $q = 2$ and denote as in [10] $\vartheta_{00} = \vartheta$, $\vartheta_{10} = \vartheta_{\frac{1}{2}, 0}$, $\vartheta_{01} = \vartheta_{0, \frac{1}{2}}$, $\vartheta_{11} = \vartheta_{\frac{1}{2}, \frac{1}{2}}$. Taking $q = 2$ in (3.38) and using the obvious equalities

$$\vartheta_{00}(-z, \tau) = \vartheta_{00}(z, \tau), \quad \vartheta_{10}(-z, \tau) = \vartheta_{10}(z, \tau), \quad \vartheta_{01}(-z, \tau) = \vartheta_{01}(z, \tau), \quad \vartheta_{11}(-z, \tau) = -\vartheta_{11}(z, \tau),$$

we obtain for all $x, u \in \mathbf{R}$

$$\vartheta_{01}(x, i)^2 \vartheta_{10}(u, i)^2 + \vartheta_{10}(x, i)^2 \vartheta_{01}(u, i)^2 + \vartheta_{11}(x, i)^2 \vartheta_{11}(u, i)^2 = \vartheta_{00}(x, i)^2 \vartheta_{00}(u, i)^2. \quad (3.39)$$

REMARK 3.4 Identity (3.39) should be compared with Riemann's theta formulae, for example with formula (A₁) in [10, p.21]

$$\begin{aligned} (\text{A}_1) \quad \vartheta_{00}(x, \tau)^2 \vartheta_{00}(u, \tau)^2 + \vartheta_{11}(x, \tau)^2 \vartheta_{11}(u, \tau)^2 &= \vartheta_{01}(x, \tau)^2 \vartheta_{01}(u, \tau)^2 + \vartheta_{10}(x, \tau)^2 \vartheta_{10}(u, \tau)^2 \\ &= \vartheta_{00}(x+u, \tau) \vartheta_{00}(x-u, \tau) \vartheta_{00}(0, \tau)^2. \end{aligned}$$

Nevertheless, it is not clear to us whether (3.39) is a mere consequence of Riemann type identities.

A Appendix

Let $\alpha \in [0, 1)$, $\rho = e^{2\pi i \alpha}$, u, v be unitaries such that $uv = \rho vu$, $A_\alpha = C^*(u, v)$ and σ be the order four automorphism of A_α defined by $\sigma(u) = v$, $\sigma(v) = u^{-1}$. Denote by E the conditional expectation from A_α onto A_α^σ defined by

$$E(x) = \frac{1}{4} (x + \sigma(x) + \sigma^2(x) + \sigma^3(x)) \quad x \in A_\alpha.$$

For any $n, m \in \mathbf{Z}$, set

$$\begin{aligned} [n, m] &= \rho^{-\frac{nm}{2}} (u^n v^m + u^{-n} v^{-m}) = 2 \rho^{-\frac{nm}{2}} (u^n v^m + \sigma^2(u^n v^m)), \\ \{n, m\} &= 4 \rho^{-\frac{nm}{2}} E(u^n v^m) = \rho^{-\frac{nm}{2}} (u^n v^m + u^{-n} v^{-m} + v^n u^{-m} + v^{-n} u^m) \\ &= \rho^{-\frac{nm}{2}} (u^n v^m + u^{-n} v^{-m}) + \rho^{\frac{nm}{2}} (u^{-m} v^n + u^m v^{-n}) = [n, m] + [-m, n]. \end{aligned}$$

The following properties of $[n, m]$ are easy to check ([3])

$$[n, m]^* = [n, m] = [-n, -m], \quad (\text{A.1})$$

$$[n, m][k, l] = \rho^{\frac{nl-mk}{2}} [n+k, m+l] + \rho^{\frac{mk-nl}{2}} [n-k, m-l]. \quad (\text{A.2})$$

Using (A1) and (A2) it is plain to check that for all $n, m, k, l \in \mathbf{Z}$

$$\{n, m\}^* = \{n, m\} = \{-m, n\} = \{-n, -m\} = \{m, -n\}, \quad (\text{A.3})$$

$$\begin{aligned} \{n, m\} \{k, l\} &= \rho^{\frac{nl-mk}{2}} \{n+k, m+l\} + \rho^{\frac{mk-nl}{2}} \{n-k, m-l\} + \rho^{\frac{nk+ml}{2}} \{n-l, m+k\} \\ &\quad + \rho^{\frac{-nk-ml}{2}} \{n+l, m-k\}. \end{aligned} \quad (\text{A.4})$$

PROPOSITION A.1 *If $\alpha \neq 0$ and $\alpha \neq \frac{1}{2}$, then $A_\alpha^\sigma = C^*(\{n, 0\}; n \geq 0)$.*

Proof. The linear span of $(u^n v^m)_{n,m \in \mathbf{Z}}$ is dense in A_α and the conditional expectation E continuous, hence $\text{span}(\{n, m\})_{n,m \in \mathbf{Z}}$ is dense in A_α^σ . Denote by B the C^* -algebra generated by $(\{n, 0\})_{n \in \mathbf{N}}$. According to (A3), it suffices to prove that for all $n, m \in \mathbf{N}$

$$\{n, m\} \in B. \quad (\text{A.5})$$

According to (A4) and (A3) we have

$$\{n, 0\} \{k, 0\} = \{n+k, 0\} + \{n-k, 0\} + \rho^{\frac{nk}{2}} \{n, k\} + \rho^{-\frac{nk}{2}} \{k, n\}, \quad (\text{A.6})$$

whence $\{n, k\} \in B$ if and only if $\{k, n\} \in B$. The previous equality gives also for all $n \in \mathbf{Z}$

$$\rho^{\frac{n}{2}} \{n, 1\} + \rho^{-\frac{n}{2}} \{1, n\} \in B. \quad (\text{A.7})$$

In particular $\{1, 1\} \in B$ and using again (A4), we get for all $n \in \mathbf{Z}$

$$\{n, 0\} \{1, 1\} = \rho^{\frac{n}{2}} \{n+1, 1\} + \rho^{-\frac{n}{2}} \{1, n-1\} + \rho^{\frac{n}{2}} \{n-1, 1\} + \rho^{-\frac{n}{2}} \{1, n+1\}. \quad (\text{A.8})$$

Let $m \geq 1$ and assume that $\{k, 1\} \in B$ for all $0 \leq k \leq m$. Then $\{1, k\} \in B$ for all $0 \leq k \leq m$ and by (A8)

$$\rho^{\frac{m}{2}} \{m+1, 1\} + \rho^{-\frac{m}{2}} \{1, m+1\} \in B. \quad (\text{A.9})$$

Taking $n = m+1$ in (A7) and using (A9) and

$$\begin{vmatrix} \rho^{\frac{m+1}{2}} & \rho^{-\frac{m+1}{2}} \\ \rho^{\frac{m}{2}} & \rho^{-\frac{m}{2}} \end{vmatrix} = \rho^{\frac{1}{2}} - \rho^{-\frac{1}{2}} \neq 0,$$

we conclude that $\{m+1, 1\} \in B$, hence $\{n, 1\} \in B$ and $\{1, n\} \in B$ for all $n \geq 0$.

Finally, we prove (A5) by induction. Assume that for some $k \geq 1$ we have

$$\{n, m\} \in B \text{ for all } n \geq 0 \text{ and } k \geq n \geq 0 \text{ or for all } k \geq n \geq 0 \text{ and } m \geq 0. \quad (\text{A.10})$$

To conclude, it will suffice to prove that

$$\{n, k+1\} \in B, \quad \text{for all } n \geq 0.$$

This holds for $n = 0, 1, \dots, k$. By (A10) and

$$\{k, k\} \{1, 1\} = \{k+1, k+1\} + \{k-1, k-1\} + \rho^k \{k-1, k+1\} + \rho^{-k} \{k+1, k-1\} \in B,$$

$$\{k+1, k\} \{1, 1\} = \rho^{\frac{1}{2}} \{k+2, k+1\} + \rho^{-\frac{1}{2}} \{k, k-1\} + \rho^{k+\frac{1}{2}} \{k, k+1\} + \rho^{-k-\frac{1}{2}} \{k+2, k-1\} \in B,$$

$$\{k+2, k\} \{1, 1\} = \rho \{k+3, k+1\} + \rho^{-1} \{k+1, k-1\} + \rho^{k+1} \{k+1, k+1\}$$

$$+ \rho^{-k-1} \{k+3, k-1\} \in B \quad \text{etc.}$$

we get $\{k+1, k+1\}, \{k+2, k+1\}, \{k+3, k+1\}, \dots \in B$.

QED

COROLLARY A.2 *If $\alpha \notin \mathbf{Q}$, then*

$$A_\alpha^\sigma = C^*(\{1, 0\}, \{2, 0\}) = C^*(u + v + u^{-1} + v^{-1}, u^2 + v^2 + u^{-2} + v^{-2}).$$

Proof. Denote by C the $*$ -algebra generated by $\{1, 0\}$ and $\{2, 0\}$. According to the previous proposition it suffices to show that $\{n, 0\} \in C$ for all $n \geq 3$. Firstly, notice that

$$\{1, 0\}^2 = \{2, 0\} + \{0, 0\} + (\rho^{\frac{1}{2}} + \rho^{-\frac{1}{2}}) \{1, 1\} \in C$$

yields $\{1, 1\} \in C$. Then, we prove by induction that

$$\{1, 0\}, \dots, \{n, 0\}, \{1, 1\}, \dots, \{n-1, 1\} \in C \quad \text{for all } n \geq 2.$$

Assume that this is the case for some $n \geq 2$. By (A6) we get

$$\rho^{\frac{n}{2}} \{n-2, 1\} + \rho^{-\frac{n}{2}} \{1, n-2\} = \{n-2, 0\} \{1, 0\} - \{n-1, 0\} - \{n-3, 0\} \in C,$$

hence $\{1, n-2\} = \{2-n, 1\} \in C$. By (A8) and the induction hypotheses

$$\rho^{\frac{n-1}{2}} \{n, 1\} + \rho^{-\frac{n-1}{2}} \{1, n\} = \{n-1, 0\} \{1, 1\} - \rho^{-\frac{n-1}{2}} \{1, n-2\} - \rho^{\frac{n-1}{2}} \{n-2, 1\} \in C. \quad (\text{A.11})$$

On the other hand

$$\{1, 1\} \{n, 0\} = \rho^{-\frac{n-1}{2}} \{n, 1\} + \rho^{\frac{n-1}{2}} \{1, n\} + \rho^{\frac{n-1}{2}} \{2-n, 1\} + \rho^{-\frac{n-1}{2}} \{n-2, 1\} \in C,$$

whence

$$\rho^{-\frac{n-1}{2}} \{n, 1\} + \rho^{\frac{n-1}{2}} \{1, n\} \in C. \quad (\text{A.12})$$

By (A11), (A12) and $\rho^m \neq 1$, $m \in \mathbf{Z}$, we get $\{n, 1\}, \{1, n\} \in C$. Moreover, using also

$$\{n, 0\} \{1, 0\} = \{n+1, 0\} + \{n-1, 0\} + \rho^{\frac{n}{2}} \{n, 1\} + \rho^{-\frac{n}{2}} \{1, n\} \in C,$$

we conclude that $\{n+1, 0\} \in C$. QED

References

- [1] J. Avron, P. H. M. v. Mouche, B. Simon, *On the measure of the spectrum for the almost Mathieu operator*, Commun. Math. Phys. 132(1990), 103–118.
- [2] J. Bellissard, *Gap labelling theorems for Schrödinger operators*, in From number theory to physics, M. Waldschmidt et al (eds), Springer 1992, p. 538–630.
- [3] O. Bratteli, G. A. Elliott, D. E. Evans, A. Kishimoto, *Non-commutative spheres I*, International J. Math. 3(1991), 139–166.
- [4] F. P. Boca, *The structure of higher-dimensional noncommutative tori and metric diophantine approximation*, J. Reine Angew. Math. 492(1997), 179–219.
- [5] F. Combes, *Crossed products and Morita equivalence*, Proc. London Math. Soc. 49(1984), 289–306.
- [6] D. E. Evans, G. A. Elliott, *The structure of the irrational rotation C^* -algebra*, Ann. of Math. 138(1993), 477–501.
- [7] C. Farsi, N. Watling, *Quartic algebras*, Canad. J. Math. 44(1992), 1167–1191.
- [8] D. R. Hofstadter, *Energy levels and wave functions of Bloch electrons in a rational or irrational magnetic field*, Phys. Rev. B 14(1976), 2239–2249.
- [9] J. Igusa, *Theta Functions*, Springer Verlag 1972.
- [10] D. Mumford, *Tata Lectures on Theta I*, Birkhäuser 1983.

- [11] M. Pimsner, D. Voiculescu, *Exact sequences for K -groups and Ext -groups of certain cross-product C^* -algebras*, J. Operator Theory 4(1980), 93–118.
- [12] M. Pimsner, D. Voiculescu, *Imbedding the irrational rotation C^* -algebra into an AF-algebra*, J. Operator Theory 4(1980), 201–210.
- [13] M. A. Rieffel, *C^* -algebras associated with irrational rotations*, Pacific J. Math. 93(1981), 415–429.
- [14] M. A. Rieffel, *Projective modules over higher-dimensional non-commutative tori*, Canad. J. Math. 40(1988), 257–338.